

Appendix

A Elements of Functional Analysis

We collect in this section a number of notions and results from the theory of linear operators in Banach spaces. We refer for instance to the book [76] for further details and proofs. In addition, in Section A.6 we give the definitions and some basic results on Sobolev spaces, for which we refer to the books [1, 2].

A.1 Bounded and Closed Operators

Consider the Banach spaces \mathcal{X} and \mathcal{Z} equipped with the norms $\|\cdot\|_{\mathcal{X}}$ and $\|\cdot\|_{\mathcal{Z}}$, respectively, and a linear map (or linear operator) $\mathbf{L} : \mathcal{Z} \rightarrow \mathcal{X}$. We denote by $\text{im } \mathbf{L}$ the range of \mathbf{L} ,

$$\text{im } \mathbf{L} = \{\mathbf{L}u \in \mathcal{X} ; u \in \mathcal{Z}\} \subset \mathcal{X},$$

by $\ker \mathbf{L}$ its kernel,

$$\ker \mathbf{L} = \{u \in \mathcal{Z} ; \mathbf{L}u = 0\} \subset \mathcal{Z},$$

and by $\mathbf{G}(\mathbf{L})$ its graph,

$$\mathbf{G}(\mathbf{L}) = \{(u, \mathbf{L}u) ; u \in \mathcal{Z}\} \subset \mathcal{Z} \times \mathcal{X}.$$

Definition A.1 (Bounded operator) A linear operator $\mathbf{L} : \mathcal{Z} \rightarrow \mathcal{X}$ is called a bounded linear operator, or simply a bounded operator, if \mathbf{L} is continuous. The set of bounded linear operators is denoted by $\mathcal{L}(\mathcal{Z}, \mathcal{X})$, and by $\mathcal{L}(\mathcal{X})$ if $\mathcal{Z} = \mathcal{X}$.

Properties A.2 [76, Chapter III, §2.2, §3.1]

- (i) For a linear operator $\mathbf{L} : \mathcal{Z} \rightarrow \mathcal{X}$ the following properties are equivalent:
- \mathbf{L} is continuous, i.e., \mathbf{L} is a bounded linear operator;
 - \mathbf{L} is continuous in 0;
 - $\sup\{\|\mathbf{L}u\|_{\mathcal{X}} ; u \in \mathcal{Z}, \|u\|_{\mathcal{Z}} = 1\} < \infty$.

(ii) For a bounded linear operator \mathbf{L} , the real number

$$\|\mathbf{L}\|_{\mathcal{L}(\mathcal{X}, \mathcal{X})} \stackrel{\text{def}}{=} \sup_{\|u\|_{\mathcal{X}}=1} \|\mathbf{L}u\|_{\mathcal{X}} = \sup_{0 < \|u\|_{\mathcal{X}} \leq 1} \frac{\|\mathbf{L}u\|_{\mathcal{X}}}{\|u\|_{\mathcal{X}}} = \sup_{\|u\|_{\mathcal{X}} \neq 0} \frac{\|\mathbf{L}u\|_{\mathcal{X}}}{\|u\|_{\mathcal{X}}}$$

is called *norm* of \mathbf{L} .

(iii) The set of bounded linear operators $\mathcal{L}(\mathcal{X}, \mathcal{X})$ is a Banach space when equipped with the norm $\|\cdot\|_{\mathcal{L}(\mathcal{X}, \mathcal{X})}$.

Definition A.3 (Closed operator) A linear operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ defined on a linear subspace $D(\mathbf{L}) \subset \mathcal{X}$ is called a closed linear operator, or simply a closed operator, if its graph $G(\mathbf{L})$ is a closed set in $\mathcal{X} \times \mathcal{X}$. The set of closed linear operators is denoted by $\mathcal{C}(\mathcal{X})$.

Properties A.4 [76, Chapter III, §5.2, Theorem 5.20]

- (i) A linear operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ is closed if and only if for any sequence $(u_n)_{n \in \mathbb{N}} \subset D(\mathbf{L})$ such that $u_n \rightarrow u$ in \mathcal{X} and $\mathbf{L}u_n \rightarrow v$ in \mathcal{X} , we have that $u \in D(\mathbf{L})$ and $\mathbf{L}u = v$.
- (ii) The sum of a closed operator with a bounded operator is a closed operator. However, the sum of two closed operators is not always a closed operator.
- (iii) A closed operator with domain $D(\mathbf{L}) = \mathcal{X}$ is bounded (closed graph theorem).
- (iv) For a closed operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ the domain $D(\mathbf{L})$ equipped with the norm

$$\|u\|_{\mathbf{L}} = (\|u\|_{\mathcal{X}}^2 + \|\mathbf{L}u\|_{\mathcal{X}}^2)^{1/2}$$

is a Banach space, and the injection $i : D(\mathbf{L}) \rightarrow \mathcal{X}$ is bounded. This norm is also called *the graph norm*.

- (v) A closed operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ belongs to $\mathcal{L}(D(\mathbf{L}), \mathcal{X})$ when $D(\mathbf{L})$ is equipped with the graph norm $\|\cdot\|_{\mathbf{L}}$.

A.2 Resolvent and Spectrum

Definition A.5 Consider a linear operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$.

(i) We call resolvent set of \mathbf{L} the set of complex numbers

$$\rho(\mathbf{L}) = \{\lambda \in \mathbb{C} ; (\lambda \mathbb{I} - \mathbf{L}) \text{ invertible and } (\lambda \mathbb{I} - \mathbf{L})^{-1} \in \mathcal{L}(\mathcal{X})\}.$$

The operator $(\lambda \mathbb{I} - \mathbf{L})^{-1}$, for $\lambda \in \rho(\mathbf{L})$, is called the resolvent of \mathbf{L} .

(ii) We call the spectrum of \mathbf{L} the complement of the resolvent set,

$$\sigma(\mathbf{L}) = \mathbb{C} \setminus \rho(\mathbf{L}).$$

- (iii) A complex number $\lambda \in \mathbb{C}$ is called an eigenvalue of \mathbf{L} if $\ker(\lambda \mathbb{I} - \mathbf{L}) \neq \{0\}$. The kernel $\ker(\lambda \mathbb{I} - \mathbf{L})$ is called the eigenspace associated with the eigenvalue

- λ , and any element $u \in \ker(\lambda\mathbb{I} - \mathbf{L}) \setminus \{0\}$ is called an eigenvector associated with the eigenvalue λ .
- (iv) For an eigenvalue $\lambda \in \sigma(\mathbf{L})$, the dimension of $\ker(\lambda\mathbb{I} - \mathbf{L})$ is called the geometric multiplicity of λ . An eigenvalue with geometric multiplicity one is called geometrically simple.
 - (v) For an isolated eigenvalue $\lambda \in \sigma(\mathbf{L})$, the dimension of the largest subspace $\mathcal{X}_\lambda \subset D(\mathbf{L})$ which is invariant under the action of \mathbf{L} and such that $\sigma(\mathbf{L}|_{\mathcal{X}_\lambda}) = \{\lambda\}$ is called the algebraic multiplicity of λ . An eigenvalue with algebraic multiplicity one is called algebraically simple or simple.
 - (vi) An eigenvalue is called semisimple if its algebraic and geometric multiplicities are the same.

Properties A.6 [76, Chapter III, §6.1, §6.3]

- (i) The spectrum of a closed operator $\mathbf{L} \in \mathcal{C}(\mathcal{X})$ is a closed set.
- (ii) The spectrum of a bounded operator $\mathbf{L} \in \mathcal{L}(\mathcal{X})$ is a closed, bounded, nonempty set.
- (iii) For a closed operator $\mathbf{L} \in \mathcal{C}(\mathcal{X})$, if $\lambda \in \rho(\mathbf{L})$, then the resolvent $(\lambda\mathbb{I} - \mathbf{L})^{-1} : \mathcal{X} \rightarrow D(\mathbf{L})$ is a bounded operator, when $D(\mathbf{L})$ is equipped with the graph norm, i.e., $(\lambda\mathbb{I} - \mathbf{L})^{-1} \in \mathcal{L}(\mathcal{X}, D(\mathbf{L}))$.
- (iv) The map $\lambda \mapsto (\lambda\mathbb{I} - \mathbf{L})^{-1}$ is holomorphic from $\rho(\mathbf{L})$ into $\mathcal{L}(\mathcal{X})$.
- (v) For $\lambda \in \rho(\mathbf{L})$, the resolvent $(\lambda\mathbb{I} - \mathbf{L})^{-1} : \mathcal{X} \rightarrow D(\mathbf{L})$ commutes with \mathbf{L} , i.e.,

$$\mathbf{L}(\lambda\mathbb{I} - \mathbf{L})^{-1}u = (\lambda\mathbb{I} - \mathbf{L})^{-1}\mathbf{L}u \text{ for all } u \in D(\mathbf{L}).$$

- (vi) For $\lambda, \mu \in \rho(\mathbf{L})$, the resolvents $(\lambda\mathbb{I} - \mathbf{L})^{-1}$ and $(\mu\mathbb{I} - \mathbf{L})^{-1}$ commute and

$$(\lambda\mathbb{I} - \mathbf{L})^{-1} - (\mu\mathbb{I} - \mathbf{L})^{-1} = (\mu - \lambda)(\lambda\mathbb{I} - \mathbf{L})^{-1}(\mu\mathbb{I} - \mathbf{L})^{-1}.$$

- (vii) For an operator \mathbf{L} , we call *the extended spectrum* the set $\sigma_\infty(\mathbf{L}) \subset \mathbb{C} \cup \{\infty\}$ defined by

$$\sigma_\infty(\mathbf{L}) = \begin{cases} \sigma(\mathbf{L}) & \text{if } \sigma(\mathbf{L}) \text{ is bounded,} \\ \sigma(\mathbf{L}) \cup \{\infty\} & \text{if } \sigma(\mathbf{L}) \text{ is unbounded.} \end{cases}$$

Then for $\lambda, \mu \in \rho(\mathbf{L})$ the spectrum of the resolvent satisfies

$$\sigma((\lambda\mathbb{I} - \mathbf{L})^{-1}) = \{(\lambda - \mu)^{-1} ; \mu \in \sigma_\infty(\mathbf{L})\}. \quad (\text{A.1})$$

Theorem A.7 (Spectral decomposition [76, Chapter III, Theorem 6.17]) Consider a closed operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$. Assume that $\sigma(\mathbf{L}) = F \cup G$, with $F \cap G = \emptyset$ and $F \subset \mathbb{C}$ a closed, bounded set, such that there exists a rectifiable, simple, closed curve Γ which encloses an open set containing F in its interior and G in its exterior. Then there exists a decomposition of $\mathcal{X} = \mathcal{X}_F \oplus \mathcal{X}_G$, with \mathcal{X}_F and \mathcal{X}_G invariant under the action of \mathbf{L} , such that the spectra of the restrictions $\mathbf{L}|_{\mathcal{X}_F}$ and $\mathbf{L}|_{\mathcal{X}_G}$ coincide with F and G , respectively. Furthermore, $\mathbf{L}|_{\mathcal{X}_F}$ is a bounded operator, $\mathbf{L}|_{\mathcal{X}_F} \in \mathcal{L}(\mathcal{X}_F)$, and the unique spectral projection $\mathbf{P}_F : \mathcal{X} \rightarrow \mathcal{X}_F$ which

commutes with \mathbf{L} is given by the Dunford integral formula

$$\mathbf{P}_F = \frac{1}{2\pi i} \int_{\Gamma} (\lambda \mathbb{I} - \mathbf{L})^{-1} d\lambda.$$

- Remark A.8** (i) The result in the above theorem still holds when the curve Γ is replaced by a finite number of rectifiable, simple, closed curves.
(ii) In the particular case when $F = \{\lambda\}$ is reduced to one point, λ is an isolated point of the spectrum. If the dimension of \mathcal{X}_F is finite, then λ is an eigenvalue of \mathbf{L} , and the dimension of \mathcal{X}_F is the algebraic multiplicity of λ [76, Chapter III §6.5].

A.3 Compact Operators and Operators with Compact Resolvent

Definition A.9 (Compact operator) A linear operator $\mathbf{L} : \mathcal{Z} \rightarrow \mathcal{X}$ is called a compact operator if for any bounded sequence $(u_n)_{n \in \mathbb{N}} \subset \mathcal{Z}$, the sequence $(\mathbf{L}u_n)_{n \in \mathbb{N}} \subset \mathcal{X}$ contains a convergent subsequence.

Properties A.10 [76, Chapter III §4.2]

- (i) A compact operator is bounded.
- (ii) The sum of two compact operators is a compact operator. For a bounded operator $\mathbf{L} : \mathcal{Y} \rightarrow \mathcal{Z}$ and a compact operator $\mathbf{K} : \mathcal{Z} \rightarrow \mathcal{X}$, the composed operator $\mathbf{K} \circ \mathbf{L} : \mathcal{Y} \rightarrow \mathcal{X}$ is compact. A similar property holds for $\mathbf{L} \circ \mathbf{K}$ (adapt the spaces).

Theorem A.11 (Spectrum [76, Chapter III, Theorem 6.26]) Consider a compact operator $\mathbf{L} : \mathcal{X} \rightarrow \mathcal{X}$. Then the following properties hold:

- (i) $0 \in \sigma(\mathbf{L})$ if \mathcal{X} is infinite-dimensional;
- (ii) any $\lambda \in \sigma(\mathbf{L})$ with $\lambda \neq 0$ is an isolated eigenvalue with finite algebraic multiplicity;
- (iii) $\sigma(\mathbf{L})$ is a countable set with at most one accumulation point in 0.

Definition A.12 (Operator with compact resolvent) A linear operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ is called an operator with compact resolvent if $\rho(\mathbf{L}) \neq \emptyset$ and for some $\lambda \in \rho(\mathbf{L})$ the operator $(\lambda \mathbb{I} - \mathbf{L})^{-1} : \mathcal{X} \rightarrow \mathcal{X}$ is compact.

Properties A.13 [76, Chapter III §6.8]

- (i) If $(\lambda \mathbb{I} - \mathbf{L})^{-1} : \mathcal{X} \rightarrow \mathcal{X}$ is a compact operator for some $\lambda \in \rho(\mathbf{L})$, then it is a compact operator for any $\lambda \in \rho(\mathbf{L})$.
- (ii) The spectrum of an operator with compact resolvent is a countable set consisting of isolated eigenvalues with finite algebraic multiplicities, with no accumulation point in \mathbb{C} .

A.4 Adjoint Operator

For a Banach space \mathcal{X} , denote by \mathcal{X}^* the dual space, i.e., the space of all continuous linear forms on \mathcal{X} , and by $\langle \cdot, \cdot \rangle$ the duality product defined by

$$\langle u, u^* \rangle = \overline{u^*(u)} \text{ for all } u \in \mathcal{X}, u^* \in \mathcal{X}^*.$$

Recall that $\mathcal{X} = \mathcal{X}^*$ when \mathcal{X} is a Hilbert space.

Definition A.14 (Adjoint operators) (i) Two linear operators $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ and $\mathbf{M} : D(\mathbf{M}) \subset \mathcal{X}^* \rightarrow \mathcal{X}^*$ are called adjoint to each other if

$$\langle \mathbf{L}u, v \rangle = \langle u, \mathbf{M}v \rangle \text{ for all } u \in D(\mathbf{L}), v \in D(\mathbf{M}).$$

(ii) If there exists a unique maximal operator \mathbf{L}^* which is adjoint to \mathbf{L} , then \mathbf{L}^* is called the adjoint of \mathbf{L} .

Properties A.15 ([76, Chapter III, §5.5, Theorem 6.22])

- (i) If $D(\mathbf{L})$ is dense in \mathcal{X} , for an operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ there is a unique adjoint operator, but this property is not true in general. The adjoint operator $\mathbf{L}^* : D(\mathbf{L}^*) \subset \mathcal{X}^* \rightarrow \mathcal{X}^*$ is constructed in the following way. The domain $D(\mathbf{L}^*)$ consists of all $v \in \mathcal{X}^*$ such that $u \mapsto \langle \mathbf{L}u, v \rangle$ is a continuous linear form on \mathcal{X} . Then there exists $w \in \mathcal{X}^*$ such that

$$\langle \mathbf{L}u, v \rangle = \langle u, w \rangle \text{ for all } u \in D(\mathbf{L}),$$

and w is unique because $D(\mathbf{L})$ is dense in \mathcal{X} . We define $\mathbf{L}^*v = w$.

- (ii) For an operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ with $D(\mathbf{L})$ dense in \mathcal{X} , the adjoint operator $\mathbf{L}^* : D(\mathbf{L}^*) \subset \mathcal{X}^* \rightarrow \mathcal{X}^*$ is closed. In addition, if \mathbf{L} is closed and the Banach space \mathcal{X} is reflexive, then \mathbf{L}^* is densely defined, i.e., $D(\mathbf{L}^*)$ is dense in \mathcal{X}^* .
- (iii) For an operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ with $D(\mathbf{L})$ dense in \mathcal{X} , we have

$$\ker \mathbf{L}^* = (\operatorname{im} \mathbf{L})^\perp \stackrel{\text{def}}{=} \{v \in \mathcal{X}^*; \langle u, v \rangle = 0 \text{ for all } u \in \operatorname{im} \mathbf{L}\}.$$

- (iv) For an operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ the resolvent set and the spectrum of the adjoint operator $\mathbf{L}^* : D(\mathbf{L}^*) \subset \mathcal{X}^* \rightarrow \mathcal{X}^*$ satisfy

$$\rho(\mathbf{L}^*) = \overline{\rho(\mathbf{L})}, \quad \sigma(\mathbf{L}^*) = \overline{\sigma(\mathbf{L})}.$$

Furthermore,

$$(\overline{\lambda \mathbb{I}} - \mathbf{L}^*)^{-1} = ((\lambda \mathbb{I} - \mathbf{L})^{-1})^* \text{ for all } \lambda \in \rho(\mathbf{L}).$$

Definition A.16 (Self-adjoint operator) A linear operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ in a Hilbert space \mathcal{X} , with domain $D(\mathbf{L})$ dense in \mathcal{X} , is called self-adjoint if its adjoint $\mathbf{L}^* : D(\mathbf{L}^*) \subset \mathcal{X} \rightarrow \mathcal{X}$ satisfies $D(\mathbf{L}) = D(\mathbf{L}^*)$ and $\mathbf{L}u = \mathbf{L}^*u$ for all $u \in D(\mathbf{L})$.

Properties A.17 [76, Chapter V, §3.4, §3.5]

- (i) The spectrum of a self-adjoint operator is real.
- (ii) The algebraic and geometric multiplicities of an isolated eigenvalue $\lambda \in \sigma(\mathbf{L})$ of a self-adjoint operator are the same, i.e., the eigenvalue is semisimple.

A.5 Fredholm Operators

Definition A.18 A bounded operator $\mathbf{L} \in \mathcal{L}(\mathcal{X}, \mathcal{X})$ is called a Fredholm operator if its kernel $\ker \mathbf{L}$ is finite-dimensional and its range $\operatorname{im} \mathbf{L}$ is closed and has finite codimension. The integer

$$\operatorname{ind}(\mathbf{L}) \stackrel{\text{def}}{=} \dim(\ker \mathbf{L}) - \operatorname{codim}(\operatorname{im} \mathbf{L}),$$

is called the Fredholm index.

Remark A.19 The above definition is easily extended to closed operators $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$, since $\mathbf{L} \in \mathcal{L}(D(\mathbf{L}), \mathcal{X})$.

Properties A.20 [76, Chapter IV, §5.1, Theorem 5.22, Theorem 5.28, §5.2]

- (i) The set of Fredholm operators is open in $\mathcal{L}(\mathcal{X}, \mathcal{X})$ and the map $\mathbf{L} \mapsto \operatorname{ind}(\mathbf{L})$ is continuous.
- (ii) For a closed operator $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$, if $\lambda \in \sigma(\mathbf{L})$ is an eigenvalue with finite algebraic multiplicity which is isolated in the spectrum of \mathbf{L} , then $\lambda \mathbb{I} - \mathbf{L}$ is a Fredholm operator with index 0.
- (iii) If $\mathbf{L} : D(\mathbf{L}) \subset \mathcal{X} \rightarrow \mathcal{X}$ is a densely defined closed operator, then \mathbf{L} is a Fredholm operator if and only if \mathbf{L}^* is a Fredholm operator. Furthermore,

$$\operatorname{ind}(\mathbf{L}^*) = -\operatorname{ind}(\mathbf{L}).$$

A.6 Basic Sobolev Spaces

We recall in this section some basic properties of the Sobolev spaces $L^2(\Omega)$ and $H^m(\Omega)$, $m \in \mathbb{N}^*$ which are used in this book. We refer to [1, Chapters II, III, V, and VI] for more general statements, and for the case of the spaces $L^p(\Omega)$ and $W^{m,p}(\Omega)$.

The Space $L^2(\Omega)$

Consider a domain $\Omega \subset \mathbb{R}^n$, and the space of complex-valued square-integrable functions on Ω

$$\mathcal{L}^2(\Omega) = \{f : \Omega \rightarrow \mathbb{C} ; f \text{ measurable and } \int_{\Omega} |f(x)|^2 dx < \infty\}.$$

This set is a linear space with respect to the natural operations (sum and multiplication by a complex number). For $f \in \mathcal{L}^2(\Omega)$, we set

$$\|f\| = \left(\int_{\Omega} |f(x)|^2 dx \right)^{1/2}. \quad (\text{A.2})$$

Then $\|\cdot\|$ is a seminorm on $\mathcal{L}^2(\Omega)$, but not a norm, since if $\|f\| = 0$, then $f = 0$ almost everywhere, only. However, $\|\cdot\|$ can be transformed into a norm by replacing $\mathcal{L}^2(\Omega)$ by the quotient with respect to the kernel of $\|\cdot\|$, i.e., by

$$L^2(\Omega) = \mathcal{L}^2(\Omega) / \ker(\|\cdot\|).$$

Clearly, the kernel of $\|\cdot\|$ consists of functions that are equal to 0 almost everywhere, so that $L^2(\Omega)$ consists of classes of functions that are equal almost everywhere. Then $\|\cdot\|$ is a norm on $L^2(\Omega)$, or, in other words $L^2(\Omega)$ is a normed space with norm $\|\cdot\|$ defined by (A.2). Furthermore, this norm corresponds to the scalar product defined by

$$\langle f, g \rangle = \int_{\Omega} f(x) \overline{g(x)} dx \text{ for all } f, g \in L^2(\Omega). \quad (\text{A.3})$$

A key property of the space $L^2(\Omega)$ is that it is complete; more precisely we have the following result.

Properties A.21 The space $L^2(\Omega)$ equipped with the scalar product $\langle \cdot, \cdot \rangle$ defined by (A.3) is a Hilbert space.

The Spaces $H^m(\Omega)$ and $H_0^m(\Omega)$

Definition A.22 Consider $m \in \mathbb{N}^*$.

(i) We define the space

$$H^m(\Omega) = \{u \in L^2(\Omega) ; D^{\alpha}u \in L^2(\Omega) \text{ for all } \alpha \in \mathbb{N}^n, |\alpha| \leq m\},$$

in which $D^{\alpha}u$ is the distributional partial derivative of u ,

$$D^{\alpha}u = \frac{\partial^{|\alpha|} u}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}},$$

for a multi-index $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ and $|\alpha| = \alpha_1 + \dots + \alpha_n$.

(ii) On $H^m(\Omega)$ we define the scalar product

$$\langle f, g \rangle_m = \sum_{|\alpha| \leq m} \langle D^{\alpha}f, D^{\alpha}g \rangle,$$

where $\langle \cdot, \cdot \rangle$ represents the scalar product in $L^2(\Omega)$, and the corresponding norm

$$\|u\|_m = \langle u, u \rangle_m^{1/2}.$$

- (iii) We define $H_0^m(\Omega)$ as the closure of $C_0^\infty(\Omega)$ in $H^m(\Omega)$, where $C_0^\infty(\Omega)$ is the space of functions of class C^∞ which have compact support in Ω .

Properties A.23 [1, Theorem 3.2]

- (i) $H^m(\Omega)$ equipped with the scalar product $\langle \cdot, \cdot \rangle_m$ is a Hilbert space.
(ii) $H^{m+j}(\Omega)$ is a dense subspace of $H^m(\Omega)$ and the imbedding $H^{m+j}(\Omega) \hookrightarrow H^m(\Omega)$ is continuous, for any $j \in \mathbb{N}^*$.

Properties A.24 (Sobolev imbedding theorem [1, Theorem 5.4]) Assume that either $\Omega = \mathbb{R}^n$ or Ω is a bounded domain in \mathbb{R}^n having a locally Lipschitz boundary, i.e., for each point x on the boundary $\partial\Omega$ of Ω there exists a neighborhood U_x such that $\partial\Omega \cap U_x$ is the graph of a Lipschitz continuous function.

- (i) For any $j \in \mathbb{N}^*$ such that $j > n/2$, we have that $H^{m+j}(\Omega) \subset C^m(\overline{\Omega})$, and the imbedding is continuous.
(ii) If Ω is an arbitrary domain in \mathbb{R}^n , the result (i) holds with $H_0^{m+j}(\Omega)$ instead of $H^{m+j}(\Omega)$.

Properties A.25 (Rellich–Kondrachov theorem [2, Theorem 3.8], [1, Theorem 6.2]) Assume that Ω is a bounded domain in \mathbb{R}^n having a locally Lipschitz boundary.

- (i) For any $j \in \mathbb{N}^*$, the imbedding $H^{m+j}(\Omega) \subset H^m(\Omega)$ is compact.
(ii) For any $j \in \mathbb{N}^*$ such that $j > n/2$, the imbedding $H^{m+j}(\Omega) \subset C^m(\overline{\Omega})$ is compact.
(iii) If Ω is an arbitrary domain in \mathbb{R}^n , the results (i) and (ii) hold with $H_0^{m+j}(\Omega)$ instead of $H^{m+j}(\Omega)$.

Spaces of Periodic Functions

An important particular case is that of spaces of periodic functions on the real line. Consider the space $L_{\text{loc}}^2(\mathbb{R})$ of measurable functions $f : \mathbb{R} \rightarrow \mathbb{C}$ satisfying $f \in L^2(a, b)$ for any bounded interval $(a, b) \subset \mathbb{R}$. We define the space of square-integrable ℓ -periodic functions by

$$L_{\text{per}}^2(0, \ell) = \{f \in L_{\text{loc}}^2(\mathbb{R}) ; f(\cdot + \ell) = f(\cdot)\},$$

and for $m \in \mathbb{N}^*$, the Sobolev spaces H^m consisting of ℓ -periodic functions by

$$H_{\text{per}}^m(0, \ell) = \{f \in L_{\text{per}}^2(\mathbb{R}) ; f^{(k)} \in L_{\text{per}}^2(\mathbb{R}) \text{ for all } k \leq m\}.$$

Notice that $L_{\text{per}}^2(0, \ell)$ can be identified with $L^2(0, \ell)$, but this is not true for the spaces H^m , $m \geq 1$. The results above, in particular the imbedding theorems, hold for these spaces, as well.

B Center Manifolds

The references in this section are to theorems, hypotheses, formulas, and remarks in Chapter 2.

B.1 Proof of Theorem 2.9 (Center Manifolds)

Consider system (2.1), and assume that Hypotheses 2.1, 2.4, and 2.7 hold. For any $u \in \mathcal{Z}$ we set

$$u = u_0 + u_h \in \mathcal{Z}, \quad u_0 = \mathbf{P}_0 u \in \mathcal{E}_0, \quad u_h = \mathbf{P}_h u \in \mathcal{Z}_h,$$

and rewrite the system (2.1) as

$$\begin{aligned} \frac{du_0}{dt} - \mathbf{L}_0 u_0 &= \mathbf{P}_0 \mathbf{R}(u) \\ \frac{du_h}{dt} - \mathbf{L}_h u_h &= \mathbf{P}_h \mathbf{R}(u). \end{aligned} \tag{B.1}$$

Modified System

We take a cut-off function $\chi : \mathcal{E}_0 \rightarrow \mathbb{R}$ of class \mathcal{C}^∞ such that

$$\chi(u_0) = \begin{cases} 1 & \text{for } \|u_0\| \leq 1 \\ 0 & \text{for } \|u_0\| \geq 2 \end{cases}, \quad \chi(u_0) \in [0, 1] \text{ for all } u_0 \in \mathcal{E}_0.$$

Since \mathcal{E}_0 is finite-dimensional such a function always exists. We use this function to modify the nonlinear terms $\mathbf{R}(u)$ outside a neighborhood of the origin, in order to be able to control the norm of the u_0 -component of the system (B.1) in the space of exponentially growing functions $\mathcal{C}_\eta(\mathbb{R}, \mathcal{E}_0)$.

We set

$$\mathbf{R}^\varepsilon(u) = \chi\left(\frac{u_0}{\varepsilon}\right) \mathbf{R}(u) \text{ for all } \varepsilon \in (0, \varepsilon_0),$$

where ε_0 is chosen such that

$$\{u = u_0 + u_h; \|u_0\|_{\mathcal{E}_0} \leq 2\varepsilon_0, \|u_h\|_{\mathcal{Z}_h} \leq \varepsilon_0\} \subset \mathcal{V},$$

with \mathcal{V} the neighborhood of the origin in Hypothesis 2.1. Then \mathbf{R}^ε is well defined in the closed set

$$\mathcal{O}_\varepsilon = \mathcal{E}_0 \times B_\varepsilon(\mathcal{Z}_h), \quad B_\varepsilon(\mathcal{Z}_h) = \{u_h \in \mathcal{Z}_h; \|u_h\| \leq \varepsilon\},$$

and satisfies

$$\mathbf{R}^\varepsilon(u) = \mathbf{R}(u) \text{ for all } u \in \mathcal{O}_\varepsilon, \|u_0\| \leq \varepsilon.$$

Consider the modified system

$$\begin{aligned}\frac{du_0}{dt} - \mathbf{L}_0 u_0 &= \mathbf{P}_0 \mathbf{R}^\varepsilon(u) \\ \frac{du_h}{dt} - \mathbf{L}_h u_h &= \mathbf{P}_h \mathbf{R}^\varepsilon(u).\end{aligned}\tag{B.2}$$

The nonlinear terms in this system now satisfy

$$\begin{aligned}\delta_0(\varepsilon) &\stackrel{\text{def}}{=} \sup_{u \in \bar{\mathcal{O}}_\varepsilon} (\|\mathbf{P}_0 \mathbf{R}^\varepsilon(u)\|_{\mathcal{E}_0}, \|\mathbf{P}_h \mathbf{R}^\varepsilon(u)\|_{\mathcal{Y}_h}) = O(\varepsilon^2) \\ \delta_1(\varepsilon) &\stackrel{\text{def}}{=} \sup_{u \in \bar{\mathcal{O}}_\varepsilon} (\|D_u \mathbf{P}_0 \mathbf{R}^\varepsilon(u)\|_{\mathcal{L}(\mathcal{Z}, \mathcal{E}_0)}, \|D_u \mathbf{P}_h \mathbf{R}^\varepsilon(u)\|_{\mathcal{L}(\mathcal{Z}, \mathcal{Y}_h)}) = O(\varepsilon).\end{aligned}\tag{B.3}$$

We prove below the existence of a “global” center manifold for this system which, due to the fact that \mathbf{R}^ε and \mathbf{R} coincide for $\|u_0\|_{\mathcal{E}_0} \leq \varepsilon$, will give the local center manifold for the system (2.1) in the theorem.

Integral Formulation

We replace system (B.2) by the integral formulation

$$\begin{aligned}u_0(t) &= \mathbf{S}_{0,\varepsilon}(u, t, u_0(0)) \stackrel{\text{def}}{=} e^{\mathbf{L}_0 t} u_0(0) + \int_0^t e^{\mathbf{L}_0(t-s)} \mathbf{P}_0 \mathbf{R}^\varepsilon(u(s)) ds \\ u_h &= \mathbf{S}_{h,\varepsilon}(u) \stackrel{\text{def}}{=} \mathbf{K}_h \mathbf{P}_h \mathbf{R}^\varepsilon(u).\end{aligned}\tag{B.4}$$

The first equation in this system is obtained by the variation of constant formula from the first equation in (B.1). Here $u_0(0) \in \mathcal{E}_0$ is arbitrary, and the exponential $e^{\mathbf{L}_0 t}$ exists since \mathcal{E}_0 is finite-dimensional. The second equation in (B.4) is obtained from Hypothesis 2.7, used with $f \in \mathcal{C}_0(\mathbb{R}, \mathcal{Y}_h)$. It is now straightforward to check that this integral system is equivalent to (B.2) for

$$u = (u_0, u_h) \in \mathcal{N}_{\eta,\varepsilon} \stackrel{\text{def}}{=} \mathcal{C}_\eta(\mathbb{R}, \mathcal{E}_0) \times \mathcal{C}_0(\mathbb{R}, \mathbf{B}_\varepsilon(\mathcal{Z}_h)),$$

with $0 < \eta \leq \gamma$ and $\varepsilon \in (0, \varepsilon_0)$. Notice that $\mathcal{N}_{\eta,\varepsilon}$ is a closed subspace of $\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}^\circ)$, so that it is complete when equipped with the norm of $\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$.

Fixed Point Argument

Our aim now is to show that (B.4) has a unique solution $u = (u_0, u_h) \in \mathcal{N}_{\eta,\varepsilon}$ for any $u_0(0) \in \mathcal{E}_0$. For this we use a fixed point argument for the map

$$\mathbf{S}_\varepsilon(u, u_0(0)) \stackrel{\text{def}}{=} (\mathbf{S}_{0,\varepsilon}(u, \cdot, u_0(0)), \mathbf{S}_{h,\varepsilon}(u)), \quad \mathbf{S}_\varepsilon(\cdot, u_0(0)) : \mathcal{N}_{\eta,\varepsilon} \rightarrow \mathcal{N}_{\eta,\varepsilon}.$$

We show that $\mathbf{S}_\varepsilon(\cdot, u_0(0))$ is well defined and that it is a contraction with respect to the norm of $\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$ for $\eta \in (0, \gamma]$, with γ the constant in Hypothesis 2.7, and ε sufficiently small.

First, Hypothesis 2.4 implies that for any $\delta > 0$ there is a constant $c_\delta > 0$ such that

$$\|e^{\mathbf{L}_0 t}\|_{\mathcal{L}(\mathcal{E}_0)} \leq c_\delta e^{\delta|t|} \text{ for all } t \in \mathbb{R}. \quad (\text{B.5})$$

Using this equality with $\delta = \eta$, we find

$$\sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \|e^{\mathbf{L}_0 t} u_0(0)\|_{\mathcal{E}_0} \right) \leq c_\eta \|u_0(0)\|_{\mathcal{E}_0},$$

which shows that the first term in $\mathbf{S}_{0,\varepsilon}(u, \cdot, u_0(0))$ belongs to $\mathcal{C}_\eta(\mathbb{R}, \mathcal{E}_0)$, for any $\eta > 0$. Next, for any $u \in \mathcal{N}_{\eta,\varepsilon}$, we have the estimates

$$\|\mathbf{P}_0 \mathbf{R}^\varepsilon(u(t))\|_{\mathcal{E}_0} \leq \delta_0(\varepsilon), \quad \|\mathbf{P}_h \mathbf{R}^\varepsilon(u(t))\|_{\mathcal{H}_h} \leq \delta_0(\varepsilon),$$

which together with (B.5) for $\delta = \eta/2$, and Hypothesis 2.7 imply

$$\begin{aligned} \sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \left\| \int_0^t e^{\mathbf{L}_0(t-s)} \mathbf{P}_0 \mathbf{R}^\varepsilon(u(s)) ds \right\|_{\mathcal{E}_0} \right) &\leq c_\delta \delta_0(\varepsilon) \sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \int_0^t e^{\delta|t-s|} ds \right) \\ &\leq \frac{2c_{\eta/2} \delta_0(\varepsilon)}{\eta}, \end{aligned}$$

and

$$\|\mathbf{K}_h \mathbf{P}_h \mathbf{R}^\varepsilon(u)\|_{\mathcal{C}_0(\mathbb{R}, \mathcal{H}_h)} \leq C(0) \delta_0(\varepsilon).$$

This shows that $\mathbf{S}_\varepsilon(u, u_0(0)) \in \mathcal{N}_{\eta,\varepsilon}$, provided $C(0) \delta_0(\varepsilon) \leq \varepsilon$, which holds for ε sufficiently small since $\delta_0(\varepsilon) = O(\varepsilon^2)$.

Now we show that the map $\mathbf{S}_\varepsilon(\cdot, u_0(0))$ is a contraction with respect to the norm of $\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$ for $\eta \in (0, \gamma]$ and sufficiently small ε . From equality (B.3) we find that

$$\begin{aligned} \|\mathbf{R}^\varepsilon(u_1) - \mathbf{R}^\varepsilon(u_2)\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Y})} &= \sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \|\mathbf{R}^\varepsilon(u_1(t)) - \mathbf{R}^\varepsilon(u_2(t))\|_{\mathcal{Y}} \right) \\ &\leq \delta_1(\varepsilon) \sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \|u_1(t) - u_2(t)\|_{\mathcal{Z}} \right) \\ &\leq \delta_1(\varepsilon) \|u_1 - u_2\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})} \end{aligned}$$

for any $u_1, u_2 \in \mathcal{N}_{\eta,\varepsilon}$. Now, using (B.5) with $\delta = \eta/2$ we obtain

$$\begin{aligned} \|\mathbf{S}_{0,\varepsilon}(u_1, \cdot, u_0(0)) - \mathbf{S}_{0,\varepsilon}(u_2, \cdot, u_0(0))\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{E}_0)} &\leq c_\delta \delta_1(\varepsilon) \sup_{t \in \mathbb{R}} \left(e^{-\eta|t|} \left| \int_0^t e^{\eta|s| + \delta|t-s|} ds \right| \right) \|u_1 - u_2\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})} \\ &\leq \frac{2c_{\eta/2} \delta_1(\varepsilon)}{\eta} \|u_1 - u_2\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})}, \end{aligned}$$

and using the estimate in Hypothesis 2.7 we find

$$\|\mathbf{S}_{h,\varepsilon}(u_1) - \mathbf{S}_{h,\varepsilon}(u_2)\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_h)} \leq C(\eta)\delta_1(\varepsilon)\|u_1 - u_2\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})}.$$

Since $\delta_1(\varepsilon) = O(\varepsilon)$ for any $\eta \in (0, \gamma]$, we can choose ε small enough such that

$$\|\mathbf{S}_\varepsilon(u_1, u_0(0)) - \mathbf{S}_\varepsilon(u_2, u_0(0))\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})} \leq \frac{1}{2}\|u_1 - u_2\|_{\mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})}.$$

Consequently, the map $\mathbf{S}_\varepsilon(\cdot, u_0(0))$ is a contraction in the complete metric space $\mathcal{N}_{\eta,\varepsilon}$.

Applying the fixed point theorem we now have the existence of a unique solution of (B.4),

$$u \stackrel{\text{def}}{=} \Phi(u_0(0)) \in \mathcal{N}_{\eta,\varepsilon}$$

for any $u_0(0) \in \mathcal{E}_0$, for any $\eta \in (0, \gamma]$, and ε sufficiently small. Clearly, this is also a solution of (B.2).

Properties of Φ

Recall that ε is chosen such that

$$C(0)\delta_0(\varepsilon) \leq \varepsilon, \quad \frac{2c_{\eta/2}\delta_1(\varepsilon)}{\eta} \leq \frac{1}{2}, \quad C(\eta)\delta_1(\varepsilon) \leq \frac{1}{2}.$$

Then the continuity on $[0, \gamma]$ of the map $\eta \rightarrow C(\eta)$ in Hypothesis 2.7 implies that for any $\tilde{\eta} \in (0, \gamma)$, we can choose $\varepsilon > 0$ such that these inequalities hold for all $\eta \in [\tilde{\eta}, \gamma]$. Consequently, for any $\tilde{\eta} \in (0, \gamma)$, there exists $\varepsilon > 0$ such that the unique fixed point $\Phi(u_0(0))$ belongs to $\mathcal{N}_{\eta,\varepsilon}$ for any $\eta \in [\tilde{\eta}, \gamma]$. This property is used later when showing that the center manifold is of class \mathcal{C}^k .

Next, notice that the map $u_0(0) \mapsto \mathbf{S}_{0,\varepsilon}(u, \cdot, u_0(0))$ is Lipschitz from \mathcal{E}_0 into $\mathcal{C}_\eta(\mathbb{R}, \mathcal{E}_0)$, so that the map $u_0(0) \mapsto \mathbf{S}_\varepsilon(u, u_0(0))$ is also Lipschitz. Consequently, Φ is a Lipschitz map. In addition, the uniqueness of the fixed point implies that

$$\Phi(0) = 0.$$

Construction of Ψ

We define now the map $\Psi : \mathcal{E}_0 \rightarrow \mathcal{Z}_h$ in the theorem, through

$$(u_0(0), \Psi(u_0(0))) \stackrel{\text{def}}{=} \Phi(u_0(0))(0) \text{ for all } u_0(0) \in \mathcal{E}_0,$$

i.e., by taking the component in \mathcal{Z}_h of the fixed point $\Phi(u_0(0))$ at $t = 0$. Since Φ is a Lipschitz map, we have that Ψ is also a Lipschitz map, and since $\Phi(0) = 0$, we have

$$\Psi(0) = 0.$$

We prove now that Ψ has the properties (i) and (ii) in the theorem.

First, we show that the manifold

$$\mathcal{M}_{\eta,\varepsilon} = \{(u_0, \Psi(u_0)) ; u_0 \in \mathcal{E}_0\}$$

is a global invariant manifold for the flow defined by (B.2). We define the shift operator Γ_s through

$$(\Gamma_s u)(t) = u(t+s) \text{ for all } t, s \in \mathbb{R}.$$

Since system (B.2) is autonomous, it is equivariant under the action Γ_s for any $s \in \mathbb{R}$, so that if u is a solution of (B.2), then $\Gamma_s u$ is also a solution of (B.2). Moreover, $\Gamma_s u \in \mathcal{N}_{\eta,\varepsilon}$ when $u \in \mathcal{N}_{\eta,\varepsilon}$.

Consider a solution u of (B.2) with $u(0) = (u_0(0), \Psi(u_0(0)))$ for some $u_0(0) \in \mathcal{E}_0$. Then $u = \Phi(u_0(0)) \in \mathcal{N}_{\eta,\varepsilon}$, and since $\Gamma_s u \in \mathcal{N}_{\eta,\varepsilon}$ is also a solution, from the uniqueness of the fixed point we conclude that

$$\Gamma_s u = \Phi(u_0(s)) \text{ for all } s \in \mathbb{R}.$$

Consequently,

$$u(s) = (u_0(s), \Psi(u_0(s))) \text{ for all } s \in \mathbb{R},$$

which shows that $\mathcal{M}_{\eta,\varepsilon}$ is globally invariant under the flow defined by (B.2). Since the system (B.1) coincides with (B.2) in

$$\mathcal{O}_\varepsilon = B_\varepsilon(\mathcal{E}_0) \times B_\varepsilon(\mathcal{Z}_h),$$

this proves part (i) of the theorem with $\mathcal{M}_0 = \mathcal{M}_{\eta,\varepsilon}$ and $\mathcal{O} = \mathcal{O}_\varepsilon$. Indeed, assume that u is a solution of (B.1) such that $u(0) \in \mathcal{M}_0 \cap \mathcal{O}$ and $u(t) \in \mathcal{O}$ for all $t \in [0, T]$. Then u satisfies (B.2) for all $t \in [0, T]$, and since $u(0) \in \mathcal{M}_{\eta,\varepsilon}$ and $\mathcal{M}_{\eta,\varepsilon}$ is an invariant manifold, we have $u(t) \in \mathcal{M}_{\eta,\varepsilon} = \mathcal{M}_0$ for all $t \in [0, T]$.

Consider now a solution u of (B.1) which belongs to $\mathcal{O} = \mathcal{O}_\varepsilon$ for all $t \in \mathbb{R}$. Then $u \in \mathcal{N}_{\eta,\varepsilon}$ and it is also a solution of (B.2). Consequently, $u = \Phi(u_0(0))$, so that $u(0) \in \mathcal{M}_{\eta,\varepsilon} = \mathcal{M}_0$, which proves part (ii) of the theorem.

Regularity of Ψ

We have proved so far that Ψ is a Lipschitz map. Notice that for this proof we have only used the fact that \mathbf{R} is of class \mathcal{C}^1 . It remains to show that Ψ is of class \mathcal{C}^k when \mathbf{R} is of class \mathcal{C}^k . For this, it is enough to prove that Φ is of class \mathcal{C}^k .

The major difficulty in proving this property comes from the fact that the *Nemitsky operator*

$$\mathbf{R}^\varepsilon : \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}) \rightarrow \mathcal{C}_\eta(\mathbb{R}, \mathcal{Y})$$

is not continuously differentiable, due to the growth of $u \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$ as $t \rightarrow \pm\infty$. The following properties of this operator are proved in [120, Lemma 3.7]:

- (i) $\mathbf{R}^\varepsilon : \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}) \rightarrow \mathcal{C}_\zeta(\mathbb{R}, \mathcal{Y})$ is continuous for any $\eta \geq 0$ and $\zeta > 0$;
- (ii) $\mathbf{R}^\varepsilon : \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}) \rightarrow \mathcal{C}_\zeta(\mathbb{R}, \mathcal{Y})$ is of class \mathcal{C}^k for any $0 \leq \eta < \zeta/k$ and $\zeta > 0$.

We point out that the k th order derivative exists for $\eta = \zeta/k$, but this derivative is continuous only if $\eta < \zeta/k$.

Following [120], the integral system (B.4) is written as

$$u = \mathbf{S}u_0(0) + \mathbf{K}\mathbf{R}^\varepsilon(u), \quad (\text{B.6})$$

with \mathbf{S} and \mathbf{K} linear maps defined by

$$(\mathbf{S}u_0(0))(t) = e^{\mathbf{L}_0 t} u_0(0),$$

and

$$(\mathbf{K}v)(t) = \int_0^t e^{\mathbf{L}_0(t-s)} \mathbf{P}_0(v(s)) ds + (\mathbf{K}_h \mathbf{P}_h(v))(t).$$

We already showed that

$$\mathbf{S} \in \mathcal{L}(\mathcal{E}_0, \mathcal{C}_{\tilde{\eta}}(\mathbb{R}, \mathcal{E}_0)), \quad \|\mathbf{S}u_0(0)\|_{\mathcal{C}_{\tilde{\eta}}(\mathbb{R}, \mathcal{E}_0)} \leq c_{\eta/2} \|u_0(0)\|_{\mathcal{E}_0},$$

and that $\mathbf{K}\mathbf{R}^\varepsilon : \mathcal{N}_{\eta, \varepsilon} \rightarrow \mathcal{N}_{\eta, \varepsilon}$ is a contraction for any $\eta \in [\tilde{\eta}, \gamma]$, when $\tilde{\eta} \in (0, \gamma)$ and ε is sufficiently small.

The idea is to consider the fixed point $u = \Phi(u_0(0)) \in \mathcal{N}_{\eta, \varepsilon} \subset \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$ of (B.4) found for $\eta \in [\tilde{\eta}, \gamma]$, with $\tilde{\eta}$ taken such that $0 < \tilde{\eta} < \gamma/k$, and to show that the map $\Phi : \mathcal{E}_0 \rightarrow \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z})$ is of class \mathcal{C}^k for all $\eta \in (k\tilde{\eta}, \gamma]$, with

$$D^p \Phi(u_0(0)) \in \mathcal{L}^p(\mathcal{E}_0, \mathcal{C}_{k\tilde{\eta}}(\mathbb{R}, \mathcal{Z})).$$

Here $\mathcal{L}^p(\mathcal{E}_0, \mathcal{C}_{k\tilde{\eta}}(\mathbb{R}, \mathcal{Z}))$ denotes the Banach space of p -linear continuous maps from \mathcal{E}_0 into $\mathcal{C}_{k\tilde{\eta}}(\mathbb{R}, \mathcal{Z})$. Several proofs of this result are available in the literature, all being quite long and technical. In particular, in [121] an abstract theorem for contractions on embedded Banach spaces is used, whereas in [120] a fiber contraction theorem due to Hirsch and Pugh [48] is used. While we refer to these works for further details, we only point out that the derivative $D\Phi(u_0(0))$ is the fixed point in $\mathcal{L}(\mathcal{E}_0, \mathcal{C}_{\tilde{\eta}}(\mathbb{R}, \mathcal{Z}))$ of the linear equation

$$D\Phi(u_0(0)) = \mathbf{S} + \mathbf{K}D_u \mathbf{R}^\varepsilon(\Phi(u_0(0)))D\Phi(u_0(0)),$$

which may be differentiated up to order k . In particular, this implies that $D\mathbf{P}_h \Phi(0) = 0$ and $D\Psi(0) = 0$, and ends the proof of Theorem 2.9. \square

B.2 Proof of Theorem 2.17 (Semilinear Case)

We prove here the first part of Theorem 2.17. The second part is an immediate consequence of this and of Theorem 2.9.

Estimates on the Resolvent of \mathbf{L}_h

First, the estimates on the resolvent (2.9) and (2.10) together with the fact that \mathbf{L}_h has no spectrum on the imaginary axis, i.e., $i\omega\mathbb{I} - \mathbf{L}_h$ is invertible for any $\omega \in \mathbb{R}$, imply that there exists a positive constant c_1 such that for any $\omega \in \mathbb{R}$, the following estimates hold:

$$\|(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h)} \leq \frac{c_1}{1 + |\omega|}, \quad \|(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h)} \leq \frac{c_1}{1 + |\omega|}, \quad (\text{B.7})$$

$$\|(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{X}_h)} \leq c_1, \quad \|(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)} \leq \frac{c_1}{(1 + |\omega|)^{1-\alpha}}. \quad (\text{B.8})$$

Next, we claim that there exist $\delta > 0$ and $M > 0$ such that any $\lambda \in \mathbb{C}$ satisfying

$$\lambda = \mu + i\omega, \quad |\mu| \leq \delta(1 + |\omega|)$$

belongs to the resolvent set of \mathbf{L}_h , and that the following estimates hold:

$$\|(\lambda\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h)} \leq \frac{M}{1 + |\lambda|}, \quad \|(\lambda\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{X}_h)} \leq M, \quad (\text{B.9})$$

$$\|(\lambda\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)} \leq \frac{M}{(1 + |\lambda|)^{1-\alpha}}. \quad (\text{B.10})$$

Indeed, we can write

$$\lambda\mathbb{I} - \mathbf{L}_h = (\mathbb{I} + \mu(i\omega\mathbb{I} - \mathbf{L}_h)^{-1})(i\omega\mathbb{I} - \mathbf{L}_h) = (i\omega\mathbb{I} - \mathbf{L}_h)(\mathbb{I} + \mu(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}),$$

and choosing $\delta = 1/2c_1$ from equalities (B.7) we find

$$\|\mu(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h)} \leq \frac{1}{2}, \quad \|\mu(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}\|_{\mathcal{L}(\mathcal{X}_h)} \leq \frac{1}{2}.$$

This shows that the operator $\mathbb{I} + \mu(i\omega\mathbb{I} - \mathbf{L}_h)^{-1}$ has a bounded inverse in both $\mathcal{L}(\mathcal{X}_h)$ and $\mathcal{L}(\mathcal{X}_h)$, so that $\lambda\mathbb{I} - \mathbf{L}_h$ is invertible. Furthermore, inequalities (B.7) and (B.8) imply the inequalities above on the norms of $(\lambda\mathbb{I} - \mathbf{L}_h)^{-1}$.

We set

$$\beta \stackrel{\text{def}}{=} \min\{|\operatorname{Re} \lambda|; \lambda \in \sigma(\mathbf{L}_h)\} \geq \delta > 0. \quad (\text{B.11})$$

(Recall that $\beta > \gamma > 0$, according to Hypothesis 2.4(i).)

Construction of \mathbf{S}_\pm

Consider the curves Γ_+ and Γ_- in \mathbb{C} defined by

$$\Gamma_+ = \{-\delta|\omega| + i\omega; \omega \in \mathbb{R}\}, \quad \Gamma_- = \{\delta|\omega| + i\omega; \omega \in \mathbb{R}\},$$

and oriented such that ω increases along Γ_+ and decreases along Γ_- . The results above imply that these two curves lie in the resolvent set of \mathbf{L}_h , and that for any λ on one of these two curves the estimates (B.9) and (B.10) hold.

For $t > 0$ we define

$$\mathbf{S}_+(t) = \frac{1}{2i\pi} \int_{\Gamma_+} e^{\lambda t} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \in \mathcal{L}(\mathcal{X}_h, \mathcal{Z}_h),$$

for which the estimates (B.9) and the dominated convergence theorem allows us to show that it is well defined and that the map $t \mapsto \mathbf{S}_+(t)$ is differentiable with

$$\frac{d^n \mathbf{S}_+(t)}{dt^n} = (\mathbf{L}_h)^n \mathbf{S}_+(t) \text{ for all } n \geq 1. \quad (\text{B.12})$$

Similarly, for $t < 0$ we set

$$\mathbf{S}_-(t) = \frac{1}{2i\pi} \int_{\Gamma_-} e^{\lambda t} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \in \mathcal{L}(\mathcal{X}_h, \mathcal{Z}_h),$$

for which we have that

$$\frac{d^n \mathbf{S}_-(t)}{dt^n} = (\mathbf{L}_h)^n \mathbf{S}_-(t) \text{ for all } n \geq 1. \quad (\text{B.13})$$

Furthermore, the commutativity property

$$\mathbf{L}_h(\lambda \mathbb{I} - \mathbf{L}_h)^{-1} = (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} \mathbf{L}_h$$

implies that

$$\begin{aligned} \mathbf{L}_h \mathbf{S}_+(t) u &= \mathbf{S}_+(t) \mathbf{L}_h u \text{ for all } u \in \mathcal{Z}_h, t > 0, \\ \mathbf{L}_h \mathbf{S}_-(t) u &= \mathbf{S}_-(t) \mathbf{L}_h u \text{ for all } u \in \mathcal{Z}_h, t < 0 \end{aligned}$$

and using the estimate (B.10) we show that for any fixed $\beta' < \beta$ and for $0 < \gamma' < \beta'$, there exists $M' > 0$ such that the estimates

$$\begin{aligned} \|\mathbf{S}_+(t)\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{Z}_h)} &\leq M'(1+t^{-\alpha})e^{-\gamma' t} \text{ for all } t > 0, \\ \|\mathbf{S}_-(t)\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{Z}_h)} &\leq M'(1+|t|^{-\alpha})e^{-\gamma' |t|} \text{ for all } t < 0, \end{aligned} \quad (\text{B.14})$$

hold. The following lemma is proved at the end of this section.

Lemma B.1 *The limits*

$$\mathbf{P}_- = \lim_{t \rightarrow 0^+} \mathbf{S}_+(t)|_{\mathcal{Y}_h} \in \mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h), \quad \mathbf{P}_+ = \lim_{t \rightarrow 0^-} \mathbf{S}_-(t)|_{\mathcal{Y}_h} \in \mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h), \quad (\text{B.15})$$

exist, and

$$(\mathbf{P}_+ + \mathbf{P}_-)u = u \text{ for all } u \in \mathcal{Y}_h. \quad (\text{B.16})$$

Checking Hypothesis 2.7

We now use the operators $\mathbf{S}_+(t)$ and $\mathbf{S}_-(t)$ above to solve the linear differential equation

$$\frac{du_h}{dt} = \mathbf{L}_h u_h + f(t). \quad (\text{B.17})$$

We show that for any $f \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Y}_h)$ with $\eta \in [0, \gamma]$ this equation has a unique solution $u_h \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_h)$ given by

$$u_h(t) = (\mathbf{K}_h f)(t) \stackrel{\text{def}}{=} \int_{-\infty}^t \mathbf{S}_+(t-s)f(s)ds - \int_t^{\infty} \mathbf{S}_-(t-s)f(s)ds, \quad (\text{B.18})$$

with the properties in Hypothesis 2.7.

We assume that γ' in (B.14) is such that $\beta > \gamma' > \gamma$. Then using these two estimates and the dominated convergence theorem it is straightforward to check that $u_h = \mathbf{K}_h f \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_h)$, and that the linear map $\mathbf{K}_h \in \mathcal{L}(\mathcal{C}_\eta(\mathbb{R}, \mathcal{Y}_h), \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_h))$, with norm satisfying the inequality in Hypothesis 2.7. Moreover using (B.12), (B.13), and Lemma B.1, we obtain that in \mathcal{X}_h the following holds

$$\begin{aligned} \frac{du_h}{dt} &= \lim_{s \rightarrow t^-} \mathbf{S}_+(t-s)f(s) + \lim_{s \rightarrow t^+} \mathbf{S}_-(t-s)f(s) \\ &\quad + \int_{-\infty}^t \mathbf{L}_h \mathbf{S}_+(t-s)f(s)ds - \int_t^{\infty} \mathbf{L}_h \mathbf{S}_-(t-s)f(s)ds \\ &= \mathbf{L}_h u_h(t) + (\mathbf{P}_+ + \mathbf{P}_-)f(t) \\ &= \mathbf{L}_h u_h(t) + f(t). \end{aligned}$$

Consequently, u_h is a solution of equation (B.17). It remains to prove the uniqueness of this solution.

Assume that $\tilde{u}_h(t) \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_h)$ is a solution of the homogeneous equation

$$\frac{d\tilde{u}_h}{dt} = \mathbf{L}_h \tilde{u}_h.$$

We show that $\tilde{u}_h = 0$. Take any $t_0 \in \mathbb{R}$, and define

$$\begin{aligned} \tilde{u}_+(t) &\stackrel{\text{def}}{=} \mathbf{S}_+(t_0 - t)\tilde{u}_h(t) \text{ for all } t < t_0, \\ \tilde{u}_-(t) &\stackrel{\text{def}}{=} \mathbf{S}_-(t_0 - t)\tilde{u}_h(t) \text{ for all } t > t_0. \end{aligned}$$

Then we have

$$\frac{d\tilde{u}_+(t)}{dt} = -\mathbf{S}_+(t_0 - t)\mathbf{L}_h\tilde{u}_h(t) + \mathbf{S}_+(t_0 - t)\frac{d\tilde{u}_h}{dt}(t) = 0$$

in \mathcal{X}_h for all $t < t_0$, hence

$$\tilde{u}_+(t) = \lim_{s \rightarrow -\infty} \tilde{u}_+(s) \text{ for all } t < t_0.$$

Using (B.21) and the continuous embedding from \mathcal{Y}_h into \mathcal{X}_h , it follows that for $\eta < \gamma' < \beta$, there is a constant $C_{\gamma'}$ such that

$$\|\mathbf{S}_+(t)\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{X}_h)} \leq C_{\gamma'} e^{-\gamma' t} \text{ for all } t > 0.$$

Consequently,

$$\|\tilde{u}_+(s)\|_{\mathcal{X}_h} \leq \|\mathbf{S}_+(t_0 - s)\|_{\mathcal{L}(\mathcal{X}_h, \mathcal{X}_h)} \|\tilde{u}_h(s)\|_{\mathcal{X}_h} \leq C_{\gamma'} e^{-\gamma'(t_0 - s)} e^{\eta|s|} \|\tilde{u}_h\|_{\mathcal{C}_\eta},$$

so that $\|\tilde{u}_+(s)\|_{\mathcal{X}_h} \rightarrow 0$ as $s \rightarrow -\infty$. This implies that $\tilde{u}_+(t) = 0$ for all $t < t_0$, and similarly we find that $\tilde{u}_-(t) = 0$ for all $t > t_0$. From the definitions of \mathbf{P}_+ and \mathbf{P}_- , and from Lemma B.1, we conclude that $\tilde{u}_h(t_0) = 0$. Since t_0 is arbitrary we have that $\tilde{u}_h = 0$, which completes the proof of Theorem 2.17. \square

Remark B.2 In the particular case when $\sigma^+ = \emptyset$, we can define the bounded projection $\mathbf{P}_- = \mathbb{I} - \mathbf{P}_0 = \mathbf{P}_h$. The estimates (B.9) imply in this case that the linear operator $\mathbf{L}_- = \mathbf{L}_h$ is the infinitesimal generator of an analytic semigroup $(\mathbf{S}_+(t))_{t \geq 0}$ in \mathcal{X}_h , which satisfies

$$\|\mathbf{S}_+(t)\|_{\mathcal{L}(\mathcal{X}_h)} \leq ce^{-\gamma' t} \text{ for all } t \geq 0.$$

This allows us to give a simpler proof of Theorem 2.17 in this case.

Proof (of Lemma B.1) For any $\eta > 0$ we define the paths Γ_+^η and Γ_-^η in \mathbb{C} by

$$\Gamma_+^\eta = \{-\delta|\omega| + i\omega; \omega \in \mathbb{R}, |\omega| \geq \delta^{-1}\eta\} \cup \{-\eta + i\omega; \omega \in \mathbb{R}, |\omega| \leq \delta^{-1}\eta\},$$

$$\Gamma_-^\eta = \{\delta|\omega| + i\omega; \omega \in \mathbb{R}, |\omega| \geq \delta^{-1}\eta\} \cup \{\eta + i\omega; \omega \in \mathbb{R}, |\omega| \leq \delta^{-1}\eta\},$$

and orient them such that ω increases along Γ_+^η , and decreases along Γ_-^η (see Figure B.1).

For any $\eta \in (0, \beta)$ we can rewrite $\mathbf{S}_\pm(t)$ as

$$\begin{aligned} \mathbf{S}_+(t) &= \frac{1}{2i\pi} \int_{\Gamma_+^\eta} e^{\lambda t} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \text{ for all } t > 0, \\ \mathbf{S}_-(t) &= \frac{1}{2i\pi} \int_{\Gamma_-^\eta} e^{\lambda t} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \text{ for all } t < 0. \end{aligned} \quad (\text{B.19})$$

Using the identity




Fig. B.1 Plot in the complex plane of the paths Γ_{\pm}^{η} (left) and $\tilde{\Gamma}_{\pm}^{\zeta}$ (right).

$$\mathbf{L}_h(\lambda \mathbb{I} - \mathbf{L}_h)^{-1} = -\mathbb{I}_{\mathcal{X}_h} + \lambda(\lambda \mathbb{I} - \mathbf{L}_h)^{-1}, \quad (\text{B.20})$$

which holds for any $\lambda \in \rho(\mathbf{L}_h)$, we obtain

$$\mathbf{S}_+(t) = \frac{1}{2i\pi} \left(\int_{\Gamma_+^{\eta}} \frac{e^{\lambda t}}{\lambda} d\lambda \right) \mathbb{I}_{\mathcal{X}_h} + \frac{1}{2i\pi} \int_{\Gamma_+^{\eta}} \frac{e^{\lambda t}}{\lambda} \mathbf{L}_h(\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda.$$

The first integral in the right hand side of this equality is independent of η , and by taking the limit as $\eta \rightarrow \infty$ we conclude that this integral vanishes.

Next, using (B.9) and the fact that $\alpha \in [0, 1)$ we find that

$$\|\mathbf{S}_+(t)\|_{\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)} \leq C_{\eta} e^{-\eta t} \text{ for all } t > 0, \quad (\text{B.21})$$

for any $\eta \in (0, \beta)$, and we conclude that

$$\mathbf{P}_- = \lim_{t \rightarrow 0^+} \mathbf{S}_+(t)|_{\mathcal{Y}_h} = \frac{1}{2i\pi} \int_{\Gamma_+^{\eta}} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \in \mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)$$

is well defined. Similarly, we find

$$\mathbf{P}_+ = \lim_{t \rightarrow 0^-} \mathbf{S}_-(t)|_{\mathcal{Y}_h} = \frac{1}{2i\pi} \int_{\Gamma_-^{\eta}} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \in \mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)$$

for any $\eta \in (0, \beta)$, which proves the first part of the lemma.

In order to prove (B.16), we define for $\zeta > 0$ the paths $\tilde{\Gamma}_{\pm}^{\zeta}$ by

$$\begin{aligned} \tilde{\Gamma}_+^{\zeta} &= \{\mu + i\delta^{-1}|\mu|; \mu \in \mathbb{R}, |\mu| \geq \delta\zeta\} \cup \{\mu + i\zeta; \mu \in \mathbb{R}, |\mu| \leq \delta\zeta\}, \\ \tilde{\Gamma}_-^{\zeta} &= \{\mu - i\delta^{-1}|\mu|; \mu \in \mathbb{R}, |\mu| \geq \delta\zeta\} \cup \{\mu - i\zeta; \mu \in \mathbb{R}, |\mu| \leq \delta\zeta\}, \end{aligned}$$

oriented such that μ decreases along $\tilde{\Gamma}_+^{\zeta}$, and increases along $\tilde{\Gamma}_-^{\zeta}$. Now observe that the operators

$$\mathbf{B}_+ \stackrel{\text{def}}{=} \frac{1}{2i\pi} \int_{\tilde{\Gamma}_+^\zeta} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda,$$

$$\mathbf{B}_- \stackrel{\text{def}}{=} \frac{1}{2i\pi} \int_{\tilde{\Gamma}_-^\zeta} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda,$$

are independent of ζ , and that the dominated convergence theorem shows their limit in $\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)$, as $\zeta \rightarrow \infty$, vanishes. Consequently, $\mathbf{B}_\pm = 0$.

Next, for $\eta = \delta\zeta$, we define the oriented clockwise rectangular path

$$\Gamma_\eta = \Gamma_+^\eta + \Gamma_-^\eta - \tilde{\Gamma}_+^\zeta - \tilde{\Gamma}_-^\zeta.$$

Then we have

$$\begin{aligned} \mathbf{P}_+ + \mathbf{P}_- &= \mathbf{B}_+ + \mathbf{B}_- + \frac{1}{2i\pi} \int_{\Gamma_\eta} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \\ &= \frac{1}{2i\pi} \int_{\Gamma_\eta} \frac{\mathbf{L}_h}{\lambda} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda \end{aligned}$$

in $\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)$, and by (B.20),

$$\mathbf{P}_+ + \mathbf{P}_- = \frac{1}{2i\pi} \int_{\Gamma_\eta} (\lambda \mathbb{I} - \mathbf{L}_h)^{-1} d\lambda - \frac{1}{2i\pi} \left(\int_{\Gamma_\eta} \frac{d\lambda}{\lambda} \right) \mathbb{I}_{\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)},$$

where $\mathbb{I}_{\mathcal{L}(\mathcal{Y}_h, \mathcal{X}_h)}$ denotes the continuous embedding from \mathcal{Y}_h into \mathcal{X}_h . The first integral on the right hand side of this equality vanishes, since the interior of the rectangle Γ_η does not contain any singularity of $(\lambda \mathbb{I} - \mathbf{L}_h)^{-1}$, which then proves (B.16), and completes the proof of the lemma. \square

B.3 Proof of Theorem 3.9 (Nonautonomous Vector Fields)

The proof of Theorem 3.9 in Chapter 2 follows exactly the proof of Theorem 2.9 given in Section B.1. The main difference is that we replace the arbitrary data $u_0(0)$ in the integral formulation (B.4) by $u_0(s) = v_0$, which modifies trivially (B.4). The fixed point is now denoted by $\Phi(v_0, s)$, so that the corresponding solution of the system (3.9), modified by the cut-off function, is

$$u(v_0, s, t) = u_0(v_0, s, t) + (\Phi(v_0, s))(t),$$

where $u_0(v_0, s, s) = v_0 \in \mathcal{E}_0$. We set

$$\Psi(u_0, t) \stackrel{\text{def}}{=} (\Phi(u_0, t))(t).$$

Then the uniqueness of the fixed point implies that

$$(\Phi(u_0(v_0, s, \tau), \tau))(t) = (\Phi(v_0, s))(t),$$

hence

$$\Psi(u_0(v_0, s, t), t) = (\Phi(v_0, s))(t).$$

This proves that

$$u_0(v_0, s, t) + \Psi(u_0(v_0, s, t), t) = u(v_0, s, t),$$

i.e., the set

$$\{(t, u_0 + \Psi(u_0, t)) ; (u_0, t) \in \mathcal{E}_0 \times \mathbb{R}\}$$

is an integral manifold for (3.9) modified by the cut-off function. Restricting to the ball $B_\varepsilon(\mathcal{E}_0)$, this implies property (i) of the theorem. Property (ii) is obtained as for Theorem 2.9. \square

We conclude this section with a brief proof of the particular cases in Corollary 3.11.

Property (i) results from a standard property of τ -periodic systems, which implies here that

$$u(v_0, s, t) = u(v_0, s + \tau, t + \tau).$$

This leads directly to

$$\Psi(u_0, t) = \Psi(u_0, t + \tau).$$

Part (ii) is obtained from the integral formulations (B.4) for both $u(v_0, s, t)$ and $u_\infty(v_0, s, t)$ by estimating, in a straightforward way, their difference as $t \rightarrow \infty$. \square

B.4 Proof of Theorem 3.13 (Equivariant Systems)

The uniqueness of the global center manifold for the modified system (B.2) in Section B.1 implies that this manifold is invariant under \mathbf{T} , provided system (B.2) is equivariant under \mathbf{T} . Since equation (2.1) is equivariant under \mathbf{T} , the modified is also equivariant when the cut-off function χ satisfies

$$\chi(\mathbf{T}_0 u_0) = \chi(u_0) \text{ for all } u_0 \in \mathcal{E}_0. \quad (\text{B.22})$$

Taking the Euclidean norm in \mathcal{E}_0 , which is finite-dimensional, for any isometry \mathbf{T}_0 on \mathcal{E}_0 we can choose χ to be a smooth function of $\|u_0\|^2$, such that (B.22) holds. Consequently, the result in the theorem follows from the fact that \mathbf{T}_0 is an isometry on \mathcal{E}_0 . \square

B.5 Proof of Theorem 3.22 (Empty Unstable Spectrum)

With the notations from Section B.1, assume that $u(\cdot; u(0)) \in \mathcal{C}^0(\mathbb{R}^+, \mathcal{Z})$ is a solution of (2.1), which belongs to $\mathcal{O}_{\varepsilon/4}$ for all $t \geq 0$. Consider $\bar{u}(t, u(0))$ defined through

$$\bar{u}(t; u(0)) = \begin{cases} u(t; u(0)) & \text{for } t \geq 0 \\ \tilde{u}(t; u(0)) & \text{for } t \leq 0, \end{cases}$$

where $\tilde{u}(\cdot; u(0))$ is the solution of the modified equation

$$\frac{d\tilde{u}}{dt} = \mathbf{L}_0 \tilde{u} + \mathbf{P}_0 \mathbf{R}^\varepsilon(\tilde{u}), \quad \tilde{u}(0) = u(0).$$

Notice that $\mathbf{P}_- \tilde{u}(t) = \mathbf{P}_- u(0)$, where $\mathbf{P}_- = \mathbb{I} - \mathbf{P}_0 = \mathbf{P}_h$. Then we find that

$$\sup_{t \leq 0} (e^{\eta t} \|\mathbf{P}_0 \bar{u}(t; u(0))\|_{\mathcal{E}_0}) < \infty,$$

and since

$$\|\mathbf{P}_0 u(t; u(0))\|_{\mathcal{E}_0} \leq \frac{\varepsilon}{4}, \quad \|\mathbf{P}_- u(t; u(0))\|_{\mathcal{Z}} \leq \frac{\varepsilon}{4} \text{ for all } t \geq 0,$$

we have that $\bar{u}(\cdot; u(0)) \in \mathcal{N}_{\eta, \varepsilon/4}$. Moreover, for all $t \in \mathbb{R}$,

$$\mathbf{P}_0 \bar{u}(t; u(0)) = e^{\mathbf{L}_0 t} \mathbf{P}_0 u(0) + \int_0^t e^{\mathbf{L}_0(t-\tau)} \mathbf{P}_0 \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0))) d\tau. \quad (\text{B.23})$$

Now assume that we have found a solution

$$z \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{E}_0) \times (\mathcal{C}_0(\mathbb{R}, B_{\varepsilon/2}(\mathcal{Z}_-)) \cap \mathcal{F}_\eta(\mathbb{R}, \mathcal{Z}_-)),$$

which means that $z(t) \rightarrow 0$ exponentially as $t \rightarrow +\infty$, of the equation

$$\begin{aligned} z(t) = & -\mathbf{P}_- \bar{u}(t; u(0)) + \mathbf{K}_h \mathbf{P}_- \mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0)) + z)(t) \\ & - \int_t^\infty e^{\mathbf{L}_0(t-\tau)} \mathbf{P}_0 [\mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)) + z(\tau)) - \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)))] d\tau. \end{aligned} \quad (\text{B.24})$$

We show below that $t \mapsto \bar{u}(t; u(0)) + z(t)$ is solution of (B.2), which belongs to $\mathcal{N}_{\eta, \varepsilon}$. As a consequence, $u(0) + z(0) \in \mathcal{M}_0$, and by construction

$$u(t, u(0) + z(0)) = u(t; u(0)) + z(t) \text{ for all } t > 0.$$

This is precisely the assertion in Theorem 3.22, since we have found a solution lying on \mathcal{M}_0 which is exponentially asymptotic to the solution $u(\cdot; u(0))$ of the initial value problem.

To end the proof, we show that $t \mapsto \bar{u}(t; u(0)) + z(t)$ is solution of (B.2). First, from (B.24) we observe that

$$\mathbf{P}_0 z(0) = - \int_0^\infty e^{-\mathbf{L}_0 \tau} \mathbf{P}_0 [\mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)) + z(\tau)) - \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)))] d\tau,$$

and using (B.23) we obtain

$$\begin{aligned} \bar{u}(t; u(0)) + z(t) &= e^{\mathbf{L}_0 t} \mathbf{P}_0(u(0) + z(0)) + (\mathbf{K}_h \mathbf{P}_- \mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0)) + z)) (t) \\ &\quad + \int_0^t e^{\mathbf{L}_0(t-\tau)} \mathbf{P}_0 \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)) + z(\tau)) d\tau. \end{aligned}$$

This is equivalent to the fact that $\bar{u}(\cdot; u(0)) + z \in \mathcal{N}_{\eta, \varepsilon}$ satisfies (B.2). It remains then to prove the existence of a solution

$$z \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{C}_0) \times (\mathcal{C}_0(\mathbb{R}, B_{\varepsilon/2}(\mathcal{Z}_-)) \cap \mathcal{F}_\eta(\mathbb{R}, \mathcal{Z}_-))$$

of (B.24).

The argument is similar to the proof of the existence of the fixed point in the proof of the center manifold theorem in Section B.1. The main difference is that the space \mathcal{C}_η is replaced by the space \mathcal{F}_η . Then for $z \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{C}_0) \times (\mathcal{C}_0(\mathbb{R}, B_{\varepsilon/2}(\mathcal{Z}_-)) \cap \mathcal{F}_\eta(\mathbb{R}, \mathcal{Z}_-))$, examining all terms in (B.24), we find that:

- (i) $\mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)) + z(\tau)) - \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0))) \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{Y})$, and

$$t \mapsto - \int_t^\infty e^{\mathbf{L}_0(t-\tau)} \mathbf{P}_0 [\mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)) + z(\tau)) - \mathbf{R}^\varepsilon(\bar{u}(\tau; u(0)))] d\tau \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{C}_0);$$

- (ii) $\mathbf{K}_h \mathbf{P}_- (\mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0)) + z) - \mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0)))) \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{Z}_-) \cap \mathcal{C}_0(\mathbb{R}, B_{\varepsilon/8}(\mathcal{Z}_-))$,
by construction and from Hypothesis 3.20, for ε sufficiently small;

- (iii) $v \stackrel{\text{def}}{=} \mathbf{K}_h \mathbf{P}_- (\mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0))) - \mathbf{P}_- \bar{u}(t; u(0))) \in \mathcal{F}_\eta(\mathbb{R}, \mathcal{Z}_-) \cap \mathcal{C}_0(\mathbb{R}, B_{3\varepsilon/8}(\mathcal{Z}_-))$.

The last property follows from the fact that $v \in \mathcal{C}_\eta(\mathbb{R}, \mathcal{Z}_-)$, and v is by construction a solution of

$$\frac{dv}{dt} = \mathbf{L}_- v$$

for $t \geq 0$, which by Hypothesis 3.20 implies the exponential convergence to 0 as $t \rightarrow \infty$, that $\mathbf{K}_h \mathbf{P}_- (\mathbf{R}^\varepsilon(\bar{u}(\cdot; u(0)))) \in \mathcal{C}_0(\mathbb{R}, B_{\varepsilon/8}(\mathcal{Z}_-))$, and that $\mathbf{P}_- \bar{u}(t; u(0)) = \mathbf{P}_- u(0) \in B_{\varepsilon/4}(\mathcal{Z}_-)$ for $t < 0$. This completes the proof of Theorem 3.22. \square

C Normal Forms

The references in this section are to theorems, hypotheses, formulas, and remarks in Chapter 3.

C.1 Proof of Lemma 1.13 (0^3 Normal Form)

The proof below can be found in [25] (see also [20] for a different proof).

We set

$$\mathbf{N}(u) = (\Phi_1(A, B, C), \Phi_2(A, B, C), \Phi_3(A, B, C)), \quad u = (A, B, C),$$

where Φ_1, Φ_2 , and Φ_3 are polynomials in (A, B, C) . Then we have $\mathbf{L}^*u = (0, A, B)$ and the characterization (1.5) leads to

$$\begin{aligned} A \frac{\partial \Phi_1}{\partial B} + B \frac{\partial \Phi_1}{\partial C} &= 0 \\ A \frac{\partial \Phi_2}{\partial B} + B \frac{\partial \Phi_2}{\partial C} &= \Phi_1 \\ A \frac{\partial \Phi_3}{\partial B} + B \frac{\partial \Phi_3}{\partial C} &= \Phi_2. \end{aligned} \tag{C.1}$$

Since A and $B^2 - 2AC$ are first integrals of the linear vector field \mathbf{L}^* , we choose the new variables

$$\tilde{A} = A, \quad \tilde{B} = B^2 - 2AC, \quad \tilde{C} = B,$$

where the change of variables is nonsingular as soon as $A \neq 0$, and define

$$\tilde{\Phi}_j(\tilde{A}, \tilde{B}, \tilde{C}) = \Phi_j(A, B, C), \quad j = 1, 2, 3.$$

Then the equations (C.1) become

$$\tilde{A} \frac{\partial \tilde{\Phi}_1}{\partial \tilde{C}} = 0, \quad \tilde{A} \frac{\partial \tilde{\Phi}_2}{\partial \tilde{C}} = \tilde{\Phi}_1, \quad \tilde{A} \frac{\partial \tilde{\Phi}_3}{\partial \tilde{C}} = \tilde{\Phi}_2, \tag{C.2}$$

so that

$$\tilde{\Phi}_1(\tilde{A}, \tilde{B}, \tilde{C}) = \phi_1(\tilde{A}, \tilde{B}).$$

We claim that ϕ_1 is a polynomial in its arguments. Indeed, by construction we have that

$$\Phi_1(A, B, C) = \Phi_1 \left(\tilde{A}, \tilde{C}, \frac{\tilde{C}^2 - \tilde{B}}{2\tilde{A}} \right),$$

which shows that ϕ_1 is a polynomial in \tilde{B} with rational coefficients in \tilde{A} . Consequently, we can write

$$\Phi_1(A, B, C) = \sum_{k \in \mathbb{N}} f_k(\tilde{A}) \tilde{B}^k = \sum_{i \in \mathbb{Z}, k \in \mathbb{N}} f_{ik} A^i \tilde{B}^k.$$

Assume that $\iota = \min\{i \in \mathbb{Z}; f_{ik} \neq 0 \text{ for some } k \in \mathbb{N}\} < 0$. Multiplying by $A^{-\iota}$ and setting $A = 0$ yields

$$\sum_{k \in \mathbb{N}} f_{\iota k} B^{2k} = 0,$$

which implies that $f_{tk} = 0$. This contradicts the assumption $\iota < 0$, so that $\iota \geq 0$. Consequently, ϕ_1 is a polynomial of (\tilde{A}, \tilde{B}) , which proves the claim.

We write now

$$\phi_1(\tilde{A}, \tilde{B}) = \tilde{A}P_1(\tilde{A}, \tilde{B}) + \psi_1(\tilde{B}),$$

where P_1 and ψ_1 are polynomials, $\psi_1(\tilde{B}) = \phi_1(0, \tilde{B})$. Solving the second equation in (C.2) leads to

$$\tilde{\Phi}_2(\tilde{A}, \tilde{B}, \tilde{C}) = BP_1(\tilde{A}, \tilde{B}) + \frac{B}{\tilde{A}}\psi_1(\tilde{B}) + \frac{\phi_2(\tilde{A}, \tilde{B})}{\tilde{A}},$$

where the same proof as above shows that ϕ_2 is a polynomial in its arguments. Multiplying this equality by \tilde{A} and setting $\tilde{A} = 0$, we obtain

$$B\psi_1(B^2) + \phi_2(0, B^2) = 0 \text{ for all } B.$$

The two terms on the right hand side of this equality have different parity, so that $\psi_1(B^2) = \phi_2(0, B^2) = 0$, and then

$$\psi_1(\tilde{B}) = 0, \quad \phi_2(0, \tilde{B}) = 0 \text{ for all } \tilde{B}.$$

Consequently, the polynomial $\phi_2(\tilde{A}, \tilde{B})$ is divisible by \tilde{A} and we can write

$$\begin{aligned} \tilde{\Phi}_1(\tilde{A}, \tilde{B}, \tilde{C}) &= AP_1(\tilde{A}, \tilde{B}), \\ \tilde{\Phi}_2(\tilde{A}, \tilde{B}, \tilde{C}) &= BP_1(\tilde{A}, \tilde{B}) + AP_2(\tilde{A}, \tilde{B}) + \psi_2(\tilde{B}), \end{aligned}$$

where P_2 and ψ_2 are polynomials.

Finally, solving the last equation in (C.2) leads to

$$\tilde{\Phi}_3(\tilde{A}, \tilde{B}, \tilde{C}) = CP_1(\tilde{A}, \tilde{B}) + BP_2(\tilde{A}, \tilde{B}) + \frac{B}{\tilde{A}}\psi_2(\tilde{B}) + \frac{\phi_3(\tilde{A}, \tilde{B})}{\tilde{A}},$$

and the same arguments as the ones above for ψ_1 and ϕ_2 apply here for ψ_2 and ϕ_3 . Then we conclude that $\psi_2(\tilde{B}) = 0$, and $\phi_3(\tilde{A}, \tilde{B})/\tilde{A} = P_3(\tilde{A}, \tilde{B})$, which is a polynomial. This completes the proof of Lemma 1.13. \square

C.2 Proof of Lemma 1.17 ($(i\omega)^2$ Normal Form)

We define

$$\mathbf{N}(u) = (\Phi_1(A, B, \bar{A}, \bar{B}), \Phi_2(A, B, \bar{A}, \bar{B}), \overline{\Phi_1(A, B, \bar{A}, \bar{B})}, \overline{\Phi_1(A, B, \bar{A}, \bar{B})}),$$

for $u = (A, B, \bar{A}, \bar{B})$, and consider the differential operator

$$\mathcal{D}^* = -i\omega A \frac{\partial}{\partial A} + (A - i\omega B) \frac{\partial}{\partial B} + i\omega \bar{A} \frac{\partial}{\partial \bar{A}} + (\bar{A} + i\omega \bar{B}) \frac{\partial}{\partial \bar{B}}.$$

Then using the characterization (1.5) in its complex form (see Remark 1.5), we find the system

$$\mathcal{D}^* \Phi_1 = -i\omega \Phi_1, \quad \mathcal{D}^* \Phi_2 = -i\omega \Phi_2 + \Phi_1.$$

First, notice that

$$\mathcal{D}^* A = -i\omega A, \quad \mathcal{D}^* B = A - i\omega B,$$

and that the equation $\mathcal{D}^* u = 0$ has the following three independent first integrals:

$$u_1 = A\bar{A}, \quad u_2 = i(A\bar{B} - \bar{A}B), \quad u_3 = i\omega \frac{B}{A} + \ln A.$$

Since $\mathcal{D}^*(\Phi_1/A) = 0$, we have that Φ_1/A is a first integral of $\mathcal{D}^* u = 0$, as well. Consequently,

$$\Phi_1(A, B, \bar{A}, \bar{B}) = A\phi(u_1, u_2, u_3) \quad (\text{C.3})$$

for some function ϕ .

We claim that ϕ is a polynomial in u_1, u_2 , and that it is independent of u_3 . Indeed, we have

$$\begin{aligned} \frac{\partial \phi}{\partial u_1} &= \frac{1}{A^2} \frac{\partial \Phi_1}{\partial \bar{A}} - \frac{B}{A^3} \frac{\partial \Phi_1}{\partial \bar{B}} \\ \frac{\partial \phi}{\partial u_2} &= \frac{-i}{A^2} \frac{\partial \Phi_1}{\partial \bar{B}} \\ \frac{\partial \phi}{\partial u_3} &= \frac{1}{i\omega} \frac{\partial \Phi_1}{\partial B} + \frac{\bar{A}}{i\omega A} \frac{\partial \Phi_1}{\partial \bar{B}}. \end{aligned}$$

Assume that Φ_1 is of degree $n-1$. Then

$$\frac{\partial^n \phi}{\partial u_j^n} = 0, \quad j = 1, 2, 3,$$

so that

$$\frac{\partial^k \phi}{\partial u_1^\alpha \partial u_2^\beta \partial u_3^\gamma} = 0 \quad \text{for } \alpha + \beta + \gamma = k \geq 3n.$$

This shows that ϕ is a polynomial in u_1, u_2, u_3 . Next, assume that ϕ depends upon u_3 . Comparing the behavior of Φ_1 and $A\phi$ in the equality (C.3), as $A \rightarrow \infty$, we obtain a contradiction between the polynomial behavior in the left hand one side and the logarithmic behavior in the right hand side. Consequently, ϕ does not depend upon u_3 and we can write

$$\Phi_1(A, B, \bar{A}, \bar{B}) = AP(u_1, u_2),$$

where P is a polynomial in its arguments.

Finally, notice that $BP(u_1, u_2)$ is a particular solution of the equation

$$\mathcal{D}^* \Phi_2 = -i\omega \Phi_2 + \Phi_1.$$

Then proceeding as above for Φ_1 , we obtain that

$$\Phi_2(A, B, \bar{A}, \bar{B}) = BP(u_1, u_2) + AQ(u_1, u_2),$$

which ends the proof of Lemma 1.17. \square

C.3 Proof of Lemma 1.18 ($0^2(i\omega)$ Normal Form)

We define

$$\mathbf{N}(u) = (\Phi_1(A, B, C, \bar{C}), \Phi_2(A, B, C, \bar{C}), \Phi_3(A, B, C, \bar{C}), \overline{\Phi_3}(A, B, C, \bar{C}))$$

for $u = (A, B, C, \bar{C})$, and consider the differential operator

$$\mathcal{D}^* = A \frac{\partial}{\partial B} - i\omega C \frac{\partial}{\partial C} + i\omega \bar{C} \frac{\partial}{\partial \bar{C}}.$$

Using characterization (1.5), we have to solve the system

$$\mathcal{D}^* \Phi_1 = 0, \quad \mathcal{D}^* \Phi_2 = \Phi_1, \quad \mathcal{D}^* \Phi_3 = -i\omega \Phi_3. \quad (\text{C.4})$$

First, notice that

$$x = A, \quad y = |C|^2, \quad z = A \ln C + i\omega B$$

are three independent first integrals of the linear equation $\mathcal{D}^* u = 0$. Using the local diffeomorphism $(A, B, C, \bar{C}) \mapsto (x, y, z, B)$, with Jacobian determinant $-A$, it is easy to show that Φ_1 expressed in the new variables, $\widetilde{\Phi}_1(x, y, z, B)$, satisfies

$$A \frac{\partial \widetilde{\Phi}_1}{\partial B} = 0.$$

Consequently, there is a function ϕ , which is smooth, except at the origin, such that

$$\Phi_1(A, B, C, \bar{C}) = \phi(x, y, z). \quad (\text{C.5})$$

We claim that ϕ is a polynomial in x, y , and that it is independent of z . Indeed, we have the equalities

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= \frac{\partial \Phi_1}{\partial A} + \frac{i \ln C}{\omega} \frac{\partial \Phi_1}{\partial B} \\ \frac{\partial \phi}{\partial y} &= \frac{1}{C} \frac{\partial \Phi_1}{\partial \bar{C}} \\ \frac{\partial \phi}{\partial z} &= \frac{1}{i\omega} \frac{\partial \Phi_1}{\partial B}, \end{aligned}$$

so that

$$\begin{aligned}\frac{\partial^n \phi}{\partial x^n} &= \frac{\partial^n \Phi_1}{\partial A^n} + \left(\frac{i \ln C}{\omega} \right)^n \frac{\partial^n \Phi_1}{\partial B^n} \\ \frac{\partial^n \phi}{\partial y^n} &= \frac{1}{C^n} \frac{\partial^n \Phi_1}{\partial \bar{C}^n} \\ \frac{\partial^n \phi}{\partial z^n} &= \frac{1}{(i\omega)^n} \frac{\partial^n \Phi_1}{\partial B^n}.\end{aligned}$$

The right hand sides of these equalities vanish for n sufficiently large, which implies that ϕ is a polynomial in its arguments. Next, assume that ϕ depends upon z . Comparing the behavior of Φ_1 and ϕ in the equality (C.5), as $A \rightarrow \infty$, we obtain a contradiction between the polynomial behavior in the left hand one side and the logarithmic behavior on the right hand side. Consequently, ϕ does not depend upon z and we can write

$$\Phi_1(A, B, C, \bar{C}) = \phi_0(A, |C|^2),$$

where ϕ_0 is a polynomial in its arguments.

Next, the second equation in (C.4) leads to

$$\Phi_2(A, B, C, \bar{C}) = \frac{B}{A} \phi_0(A, |C|^2) + \phi_1(A, |C|^2, z),$$

and using the fact that $A\Phi_2$ and ϕ_0 are polynomials, it follows that $A\phi_1 = \psi(A, |C|^2)$, with ψ a polynomial satisfying

$$B\phi_0(0, |C|^2) = \psi(0, |C|^2).$$

This implies that $\phi_0(0, |C|^2) = 0$, and that ϕ_1 is a polynomial in A , and $|C|^2$. Summarizing, there are two polynomials P_0 and P_1 in A and $|C|^2$, such that

$$\begin{aligned}\Phi_1(A, B, C, \bar{C}) &= AP_0(A, |C|^2), \\ \Phi_2(A, B, C, \bar{C}) &= BP_0(A, |C|^2) + P_1(A, |C|^2).\end{aligned}$$

Finally, the equation for Φ_3 in (C.4) leads to

$$\mathcal{D}^*(\bar{C}\Phi_3) = 0.$$

Consequently,

$$\bar{C}\Phi_3(A, B, C, \bar{C}) = \phi_2(A, |C|^2),$$

where ϕ_2 is a polynomial such that $\phi_2(A, 0) = 0$, and we conclude that there is a polynomial P_2 such that

$$\Phi_3(A, B, C, \bar{C}) = CP_2(A, |C|^2).$$

This completes the proof of Lemma 1.18. □

C.4 Proof of Lemma 1.19 (0^20^2 Normal Form)

We define

$$\mathbf{N}(u) = (\Phi_1(A, B, C, D), \Phi_2(A, B, C, D), \Phi_3(A, B, C, D), \Phi_4(A, B, C, D))$$

for $u = (A, B, C, D)$, and consider the differential operator

$$\mathcal{D}^* = A \frac{\partial}{\partial B} + C \frac{\partial}{\partial D}.$$

Using characterization (1.5), we obtain that

$$\mathcal{D}^* \Phi_1 = 0, \quad \mathcal{D}^* \Phi_2 = \Phi_1, \quad \mathcal{D}^* \Phi_3 = 0, \quad \mathcal{D}^* \Phi_4 = \Phi_3. \quad (\text{C.6})$$

First, notice that

$$A, \quad C, \quad \tilde{B} = BC - AD$$

are three independent first integrals of the equation $\mathcal{D}^* u = 0$. Then it is not difficult to show that there is a function ϕ_1 that is smooth, except at the origin, such that

$$\Phi_1(A, B, C, D) = \phi_1(A, C, \tilde{B}).$$

Furthermore, we have the identities

$$\begin{aligned} \frac{\partial^n \phi_1}{\partial A^n} &= \left(\frac{\partial}{\partial A} + \frac{D}{C} \frac{\partial}{\partial B} \right)^n \phi_1 \\ \frac{\partial^n \phi_1}{\partial C^n} &= \left(\frac{\partial}{\partial C} + \frac{B}{A} \frac{\partial}{\partial D} \right)^n \phi_1 \\ \frac{\partial^n \phi_1}{\partial \tilde{B}^n} &= \left(\frac{1}{C} \frac{\partial}{\partial B} \right)^n \phi_1, \end{aligned}$$

and the right hand sides of these equalities vanish for n sufficiently large, since Φ_1 is a polynomial. Consequently, ϕ_1 is a polynomial in its arguments.

We decompose ϕ_1 as

$$\phi_1(A, C, \tilde{B}) = A Q_1(A, C, \tilde{B}) + C Q_2(A, C, \tilde{B}) + Q_3(\tilde{B}),$$

where Q_j are polynomials in their arguments. Notice that this decomposition is not unique. Now we set

$$\Phi_2(A, B, C, D) = B Q_1(A, C, \tilde{B}) + D Q_2(A, C, \tilde{B}) + \frac{B}{A} Q_3(\tilde{B}) + \frac{1}{A} \tilde{\Phi}_2(A, B, C, D) \quad (\text{C.7})$$

so that the second equation in (C.6) leads to

$$\mathcal{D}^* \tilde{\Phi}_2 = 0,$$

where $\tilde{\Phi}_2$ is a polynomial in its arguments. The arguments used for ϕ_1 above, imply that

$$\tilde{\Phi}_2(A, B, C, D) = \phi_2(A, C, \tilde{B}),$$

with ϕ_2 polynomial in its arguments. Now multiplying (C.7) by A and taking $A = 0$ gives

$$BQ_3(BC) + \phi_2(0, C, BC)$$

for any $(B, C) \in \mathbb{R}^2$. This proves that

$$Q_3 = 0, \quad \phi_2(0, C, BC) = 0.$$

As a consequence, since B and BC are independent variables, the polynomial $\phi_2(A, C, \tilde{B})$ is divisible by A , and there is a polynomial Q_4 such that

$$\Phi_2(A, B, C, D) = BQ_1(A, C, \tilde{B}) + DQ_2(A, C, \tilde{B}) + Q_4(A, C, \tilde{B}).$$

Finally, notice that we can write

$$\begin{aligned} Q_4(A, C, \tilde{B}) &= P_3(A, C) + \tilde{B}Q_5(A, C, \tilde{B}) \\ &= P_3(A, C) + BCQ_5(A, C, \tilde{B}) - DAQ_5(A, C, \tilde{B}), \end{aligned}$$

hence we obtain the final form

$$\begin{aligned} \Phi_1(A, B, C, D) &= AP_1(A, C, \tilde{B}) + CP_2(A, C, \tilde{B}) \\ \Phi_2(A, B, C, D) &= BP_1(A, C, \tilde{B}) + DP_2(A, C, \tilde{B}) + P_3(A, C), \end{aligned} \quad (\text{C.8})$$

where

$$P_1 = Q_1 + CQ_5, \quad P_2 = Q_2 - AQ_5.$$

In the same way, from the last two equalities in (C.6) we obtain

$$\begin{aligned} \Phi_3(A, B, C, D) &= AP_4(A, C, \tilde{B}) + CP_5(A, C, \tilde{B}) \\ \Phi_4(A, B, C, D) &= BP_4(A, C, \tilde{B}) + DP_5(A, C, \tilde{B}) + P_6(A, C), \end{aligned} \quad (\text{C.9})$$

which completes the proof of Lemma 1.19. \square

C.5 Proof of Theorem 2.2 (Perturbed Normal Forms)

We have to determine two polynomials Φ_μ and \mathbf{N}_μ of degree p in \mathbb{R}^n with coefficients depending upon μ , which satisfy the equality

$$\mathcal{F}(\Phi_\mu, \mathbf{N}_\mu, \mu) = 0. \quad (\text{C.10})$$

With the notations from Hypothesis 2.1, the map $\mathcal{F} : \mathcal{H}^2 \times \mathcal{V}_\mu \rightarrow \mathcal{H}$ defined by

$$(\mathcal{F}(\Phi, \mathbf{N}, \mu))(v) \stackrel{\text{def}}{=} (\mathcal{A}_{\mathbf{L}}\Phi)(v) + \mathbf{N}(v) + \Pi_p(D\Phi(v)\mathbf{N}(v) - \mathbf{R}(v + \Phi(v), \mu)) \quad (\text{C.11})$$

is of class \mathcal{C}^k , where we recall that \mathcal{H} is the space of polynomials of degree p , and Π_p the linear map that associates to a map of class \mathcal{C}^p the polynomial of degree p in its Taylor expansion. For notational simplicity, we suppress the indices μ and write Φ and \mathbf{N} instead of Φ_μ and \mathbf{N}_μ , respectively.

For $\mu = 0$ we recover exactly the situation treated in Theorem 1.2. This means that we have a solution $(\Phi, \mathbf{N}) = (\Phi^{(0)}, \mathbf{N}^{(0)})$ of (C.10) for $\mu = 0$,

$$\mathcal{F}(\Phi^{(0)}, \mathbf{N}^{(0)}, 0) = 0,$$

which is unique when we restrict to

$$\Phi \in (\ker \mathcal{A}_{\mathbf{L}})^\perp, \quad \mathbf{N} \in \ker(\mathcal{A}_{\mathbf{L}}^*). \quad (\text{C.12})$$

In order to determine the polynomials Φ and \mathbf{N} for μ close to zero, we use the implicit function theorem to solve (C.10), together with (C.12), i.e., with

$$\mathcal{F} : (\ker \mathcal{A}_{\mathbf{L}})^\perp \times \ker(\mathcal{A}_{\mathbf{L}}^*) \times \mathcal{V}_\mu \rightarrow \mathcal{H}.$$

First, we compute the differential

$$\mathcal{D}_0 \stackrel{\text{def}}{=} D_{(\Phi, \mathbf{N})} \mathcal{F}(\Phi^{(0)}, \mathbf{N}^{(0)}, 0) : (\ker \mathcal{A}_{\mathbf{L}})^\perp \times \ker(\mathcal{A}_{\mathbf{L}}^*) \rightarrow \mathcal{H}$$

of \mathcal{F} with respect to (Φ, \mathbf{N}) at $(\Phi^{(0)}, \mathbf{N}^{(0)}, 0)$. A direct calculation gives

$$\begin{aligned} (\mathcal{D}_0(\Psi, \mathbf{M}))(v) &= (\mathcal{A}_{\mathbf{L}}\Psi)(v) + \mathbf{M}(v) + \Pi_p \left(D\Psi(v)\mathbf{N}^{(0)}(v) \right. \\ &\quad \left. + D\Phi^{(0)}(v)\mathbf{M}(v) - D\mathbf{R}(v + \Phi^{(0)}(v), 0)\Psi(v) \right), \quad (\text{C.13}) \end{aligned}$$

and we prove now that this linear map is invertible.

Denote by π_q the linear map on \mathcal{H} , which associates to a polynomial \mathbf{P} the polynomial \mathbf{P}_q obtained by suppressing the monomials in \mathbf{P} of degree different of q . With this notation, according to Theorem 1.2 we have that

$$(\Phi_0^{(0)}, \mathbf{N}_0^{(0)}) = (\Phi_1^{(0)}, \mathbf{N}_1^{(0)}) = 0,$$

since the polynomials $(\Phi^{(0)}, \mathbf{N}^{(0)})$ are at least quadratic. Identifying the homogeneous polynomials of degrees $0, \dots, p$ on the right hand side of (C.13) we obtain, successively,

$$\begin{aligned} &(\mathcal{A}_{\mathbf{L}}\Psi_0)(v) + \mathbf{M}_0(v), \\ &(\mathcal{A}_{\mathbf{L}}\Psi_1)(v) + \mathbf{M}_1(v) + D\Phi_2^{(0)}(v)\mathbf{M}_0 - D\mathbf{R}_2(v)\Psi_0, \end{aligned}$$

$$\begin{aligned}
& (\mathcal{A}_{\mathbf{L}}\Psi_2)(v) + \mathbf{M}_2(v) + D\Phi_3^{(0)}(v)\mathbf{M}_0 + D\Phi_2^{(0)}(v)\mathbf{M}_1(v) - D\mathbf{R}_2(v)\Psi_1(v) \\
& \quad - D^2\mathbf{R}_2(\Psi_0, \Phi_2^{(0)}(v)) - D\mathbf{R}_3(v)\Psi_0, \\
& (\mathcal{A}_{\mathbf{L}}\Psi_p)(v) + \mathbf{M}_p(v) + \sum_{1 \leq l \leq p} D\Phi_{l+1}^{(0)}(v)\mathbf{M}_{p-l}(v) + \sum_{1 \leq l \leq p-1} D\Psi_{p-l}(v)\mathbf{N}_{l+1}^{(0)}(v) \\
& \quad - \sum_{2 \leq q \leq p} \pi_p \left(D\mathbf{R}_q(v + \Phi^{(0)}(v))\Psi(v) \right),
\end{aligned}$$

where $\mathbf{R}_s(u) = D_u^s \mathbf{R}(0,0)(u^{(s)})/s!$. Now notice that for any degree q between 0 and p , the formulas above are of the form

$$\mathcal{A}_{\mathbf{L}}\Psi_q + \mathbf{M}_q + \mathbf{G}_q,$$

with \mathbf{G}_q depending only upon \mathbf{M}_j and Ψ_j with $j = 0, \dots, q-1$. We have seen in the proof of Theorem 1.2 that the equation

$$\mathcal{A}_{\mathbf{L}}\Psi + \mathbf{M} = \mathbf{Q}$$

has a unique solution

$$\mathbf{M} = \mathbf{P}_{\ker(\mathcal{A}_{\mathbf{L}}^*)} \mathbf{Q} \in \ker(\mathcal{A}_{\mathbf{L}}^*), \quad \Psi \in (\ker \mathcal{A}_{\mathbf{L}})^\perp,$$

for any homogeneous polynomial \mathbf{Q} of degree q . This implies that the differential \mathcal{D}_0 is invertible, and by the implicit function theorem we conclude that (C.10) has a unique solution $(\Phi_\mu, \mathbf{N}_\mu)$ satisfying (C.12). Furthermore the map $\mu \mapsto (\Phi_\mu, \mathbf{N}_\mu)$ is of class \mathcal{C}^k , which implies that the coefficients of monomials of degree q are functions of μ of class \mathcal{C}^{k-q} . This completes the proof of Theorem 2.2. \square

Remark C.1 *In the case where $\mathbf{R}(\cdot, \mu)$ is linear, equation (C.10) is affine in Φ , and we can look directly for a solution $(\Phi, \mathbf{N}) \in (\mathcal{L}(\mathbb{R}^n))^2$, i.e., in the space of polynomials of degree 1.*

D Reversible Bifurcations

The references in this section are to theorems, hypotheses, formulas, and remarks in Chapter 4.

D.1 0^{3+} Normal Form in Infinite Dimensions

We show below how to compute the principal coefficients in the normal form (2.9) when starting from an infinite-dimensional system

$$\frac{du}{dt} = \mathbf{L}u + \mathbf{R}(u, \mu), \quad (\text{D.1})$$

just like the system (4.1) in Chapter 3. We assume that the parameter μ is real, and that the system (D.1) possesses a reversibility symmetry \mathbf{S} and satisfies the hypotheses of Theorems 3.3 and 3.15 in Chapter 2. We further assume that the spectrum of the linear operator \mathbf{L} is such that $\sigma_0 = \{0\}$, where 0 is an algebraically triple and geometrically simple eigenvalue, with a symmetric eigenvector ζ_0 such that

$$\mathbf{S}\zeta_0 = \zeta_0.$$

Then the three-dimensional reduced system satisfies the hypotheses of Lemma 2.4, so that its normal form is given by (2.9).

For the computation of the coefficients in the expansion of P we proceed as in Section 3.4, and as for the reversible bifurcations in Section 4.1. We start from the equality (4.3) in Chapter 3, in which we take $v_0 = A\zeta_0 + B\zeta_1 + C\zeta_2$, and then write

$$u = A\zeta_0 + B\zeta_1 + C\zeta_2 + \tilde{\Psi}(A, B, C, \mu), \quad (\text{D.2})$$

where $\tilde{\Psi}$ takes values in \mathcal{V} . With the notations from Section 3.2.3, we consider the Taylor expansions of \mathbf{R} in (1.15) and of $\tilde{\Psi}$,

$$\tilde{\Psi}(A, B, C, \mu) = \sum_{1 \leq r+s+q+n \leq p} A^r B^s C^q \mu^n \Psi_{rsqn}, \quad \Psi_{rsq0} = 0 \text{ for } r+s+q = 1.$$

Using the reversibility symmetry we find that

$$\mathbf{R}_{r,q}((\mathbf{S}u)^{(r)}) = -\mathbf{S}\mathbf{R}_{r,q}(u^{(r)}), \quad \mathbf{S}\Psi_{rsqn} = (-1)^s \Psi_{rsqn}.$$

Now we identify the different powers of (A, B, C, μ) in the identity (4.4) in Chapter 3, which is here

$$\begin{aligned} & (\partial_A \tilde{\Psi})B + (\partial_B \tilde{\Psi})C + (\zeta_1 + \partial_B \tilde{\Psi})AP(A, \tilde{B}, \mu) + (\zeta_2 + \partial_C \tilde{\Psi})BP(A, \tilde{B}, \mu) \\ &= \mathbf{L}\tilde{\Psi} + \mathbf{R}(A\zeta_0 + B\zeta_1 + C\zeta_2 + \tilde{\Psi}, \mu). \end{aligned}$$

This leads to the following equalities, found successively at orders $\mu, A\mu, B\mu, C\mu$:

$$0 = \mathbf{L}\Psi_{0001} + \mathbf{R}_{0,1}, \quad (\text{D.3})$$

$$a\zeta_1 = \mathbf{L}\Psi_{1001} + \mathbf{R}_{1,1}\zeta_0 + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0001}), \quad (\text{D.4})$$

$$a\zeta_2 + \Psi_{1001} = \mathbf{L}\Psi_{0101} + \mathbf{R}_{1,1}\zeta_1 + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0001}), \quad (\text{D.5})$$

$$\Psi_{0101} = \mathbf{L}\Psi_{0011} + \mathbf{R}_{1,1}\zeta_2 + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{0001}); \quad (\text{D.6})$$

at orders A^2, AB, AC, BC, C^2 :

$$b\zeta_1 = \mathbf{L}\Psi_{2000} + \mathbf{R}_{2,0}(\zeta_0, \zeta_0), \quad (\text{D.7})$$

$$b\zeta_2 + 2\Psi_{2000} = \mathbf{L}\Psi_{1100} + 2\mathbf{R}_{2,0}(\zeta_0, \zeta_1), \quad (\text{D.8})$$

$$\Psi_{1100} = \mathbf{L}\Psi_{0200} + \mathbf{R}_{2,0}(\zeta_1, \zeta_1), \quad (\text{D.9})$$

$$\Psi_{1100} = \mathbf{L}\Psi_{1010} + 2\mathbf{R}_{2,0}(\zeta_0, \zeta_2), \quad (\text{D.10})$$

$$2\Psi_{0200} + \Psi_{1010} = \mathbf{L}\Psi_{0110} + 2\mathbf{R}_{2,0}(\zeta_1, \zeta_2), \quad (\text{D.11})$$

$$\Psi_{0110} = \mathbf{L}\Psi_{0020} + \mathbf{R}_{2,0}(\zeta_2, \zeta_2); \quad (\text{D.12})$$

at orders $A^3, A^2B, AB^2, B^3, A^2C, ABC$:

$$\begin{aligned} d\zeta_1 + b\Psi_{1100} &= \mathbf{L}\Psi_{3000} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{2000}) \\ &\quad + \mathbf{R}_{3,0}(\zeta_0, \zeta_0, \zeta_0), \end{aligned} \quad (\text{D.13})$$

$$\begin{aligned} d\zeta_2 + 3\Psi_{3000} + 2b\Psi_{0200} + b\Psi_{1010} &= \mathbf{L}\Psi_{2100} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{1100}) \\ &\quad + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{2000}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \zeta_1), \end{aligned} \quad (\text{D.14})$$

$$\begin{aligned} c\zeta_1 + 2\Psi_{2100} + b\Psi_{0110} &= \mathbf{L}\Psi_{1200} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0200}) \\ &\quad + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{1100}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \zeta_1), \end{aligned} \quad (\text{D.15})$$

$$\begin{aligned} c\zeta_2 + 2\Psi_{1200} &= \mathbf{L}\Psi_{0300} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0200}) \\ &\quad + \mathbf{R}_{3,0}(\zeta_1, \zeta_1, \zeta_1), \end{aligned} \quad (\text{D.16})$$

$$\begin{aligned} -2c\zeta_1 + \Psi_{2100} + b\Psi_{0110} &= \mathbf{L}\Psi_{2010} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{1010}) \\ &\quad + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{2000}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \zeta_2), \end{aligned} \quad (\text{D.17})$$

$$\begin{aligned} -2c\zeta_2 + 2\Psi_{2010} + 2\Psi_{1200} + 2b\Psi_{0020} &= \mathbf{L}\Psi_{1110} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0110}) \\ &\quad + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{1010}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{1100}) \\ &\quad + 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \zeta_2); \end{aligned} \quad (\text{D.18})$$

and at orders AC^2, B^2C, BC^2, C^3 :

$$\begin{aligned} \Psi_{1110} &= \mathbf{L}\Psi_{1020} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0020}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{1010}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_2, \zeta_2), \end{aligned} \quad (\text{D.19})$$

$$\begin{aligned} \Psi_{1110} + 3\Psi_{0300} &= \mathbf{L}\Psi_{0210} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0110}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{0200}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \zeta_2), \end{aligned} \quad (\text{D.20})$$

$$\begin{aligned} 2\Psi_{0210} &= \mathbf{L}\Psi_{0120} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0020}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{0110}) \\ &\quad + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_2, \zeta_2), \end{aligned} \quad (\text{D.21})$$

$$\Psi_{0120} = \mathbf{L}\Psi_{0030} + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{0020}) + \mathbf{R}_{3,0}(\zeta_2, \zeta_2, \zeta_2). \quad (\text{D.22})$$

Notice that due to the symmetry properties of $\mathbf{R}_{p,q}$, ζ_0 , ζ_1 , ζ_2 , and since $\mathbf{S}\Psi_{rsqn} = (-1)^s\Psi_{rsqn}$, we have that

$$\mathbf{S}\mathbf{R}_{0,1} = -\mathbf{R}_{0,1}, \quad \mathbf{S}\mathbf{R}_{1,1}\zeta_j = (-1)^{j+1}\mathbf{R}_{1,1}\zeta_j, \quad j = 0, 1, 2,$$

and

$$\begin{aligned}\mathbf{SR}_{2,0}(\zeta_j, \Psi_{rsqn}) &= (-1)^{j+s+1} \mathbf{R}_{2,0}(\zeta_j, \Psi_{rsqn}), \quad j = 0, 1, 2, \\ \mathbf{SR}_{3,0}(\zeta_j, \zeta_k, \zeta_l) &= (-1)^{j+k+l+1} \mathbf{R}_{3,0}(\zeta_j, \zeta_k, \zeta_l), \quad j, k, l \in \{0, 1, 2\}.\end{aligned}$$

The solvability conditions for these equations are now obtained by taking the duality product of each equation with the vector ζ_2^* orthogonal to the range of \mathbf{L} . Proceeding as for the other examples, we find that this vector is given by

$$\zeta_2^* = \mathbf{P}_0^* \zeta_{02}^* \in \mathcal{X}^*,$$

where \mathbf{P}_0^* is the adjoint of the projection \mathbf{P}_0 onto the three-dimensional space \mathcal{E}_0 , and ζ_{02}^* is the eigenvector associated with the eigenvalue 0 of the adjoint of \mathbf{L}_0 , the restriction of \mathbf{L} to \mathcal{E}_0 , satisfying $\langle \zeta_2, \zeta_2^* \rangle = 1$. In addition, we have that

$$\langle \zeta_0, \zeta_2^* \rangle = 0, \quad \langle \zeta_1, \zeta_2^* \rangle = 0, \quad \langle \zeta_2, \zeta_2^* \rangle = 1,$$

and since $\mathbf{S}\zeta_2 = \zeta_2$, that

$$\mathbf{S}^* \zeta_2^* = \zeta_2^*.$$

We can now solve the system (D.3)–(D.22). Since any antisymmetric vector of \mathcal{X} lies in the range of \mathbf{L} , it is straightforward to check that there is no solvability condition for the equations (D.3), (D.4), (D.6), (D.7), (D.9), (D.10), (D.12), and (D.13), (D.15), (D.17), (D.19), (D.20), (D.22). The solvability conditions for the remaining equations allow us to determine the coefficients a , c , and d .

First, the equation (D.3) gives Ψ_{0001} , defined up to an arbitrary multiple of ζ_0 . Then from (D.4) we obtain

$$\Psi_{1001} = \tilde{\Psi}_{1001} + a\zeta_2,$$

where

$$\mathbf{L}\tilde{\Psi}_{1001} + \mathbf{R}_{1,1}\zeta_0 + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0001}) = 0.$$

For (D.5) we find the solvability condition

$$2a = \langle \mathbf{R}_{1,1}\zeta_1 + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0001}) - \tilde{\Psi}_{1001}, \zeta_2^* \rangle,$$

which gives a , and from this equation we can determine Ψ_{0101} up to an arbitrary multiple of ζ_0 . From (D.6) we can find Ψ_{0011} up to an arbitrary multiple of ζ_0 , again.

Next, equation (D.7) gives

$$\Psi_{2000} = \tilde{\Psi}_{2000} + b\zeta_2 + \psi_{2000}\zeta_0,$$

where

$$\mathbf{L}\tilde{\Psi}_{2000} + \mathbf{R}_{2,0}(\zeta_0, \zeta_0) = 0.$$

For (D.8) we find the solvability condition

$$3b = \langle 2\mathbf{R}_{2,0}(\zeta_0, \zeta_1) - 2\tilde{\Psi}_{2000}, \zeta_2^* \rangle,$$

which gives b , and we can determine

$$\Psi_{1100} = \tilde{\Psi}_{1100} + 2\psi_{2000}\zeta_1,$$

where

$$\mathbf{L}\tilde{\Psi}_{1100} + 2\mathbf{R}_{2,0}(\zeta_0, \zeta_1) = 2\tilde{\Psi}_{2000}.$$

Similarly, from (D.9) and (D.10) we obtain,

$$\Psi_{0200} = \tilde{\Psi}_{0200} + 2\psi_{2000}\zeta_2, \quad \Psi_{1010} = \tilde{\Psi}_{1010} + 2\psi_{2000}\zeta_2,$$

where

$$\mathbf{L}\tilde{\Psi}_{0200} + \mathbf{R}_{2,0}(\zeta_1, \zeta_1) = \tilde{\Psi}_{1100}, \quad \mathbf{L}\tilde{\Psi}_{1010} + 2\mathbf{R}_{2,0}(\zeta_0, \zeta_2) = \tilde{\Psi}_{1100}.$$

Now for (D.11) we find the solvability condition

$$6\psi_{2000} = \langle 2\mathbf{R}_{2,0}(\zeta_1, \zeta_2) - 2\tilde{\Psi}_{0200} - \tilde{\Psi}_{1010}, \zeta_2^* \rangle,$$

and from (D.11) and (D.12) we can determine Ψ_{0110} and Ψ_{0020} .

Finally, from (D.13) and (D.14) we find

$$\Psi_{3000} = \tilde{\Psi}_{3000} + d\zeta_2 + \psi_{3000}\zeta_0, \quad \Psi_{2100} = \tilde{\Psi}_{2100} + 3\psi_{3000}\zeta_1,$$

and the coefficient d ,

$$\begin{aligned} 4d = & \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{1100}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{2000}) + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0\zeta_1), \zeta_2^* \rangle \\ & - \langle 3\tilde{\Psi}_{3000} + 2b\Psi_{0200} + b\Psi_{1010}, \zeta_2^* \rangle. \end{aligned} \quad (\text{D.23})$$

Now (D.15) and (D.16) give

$$\Psi_{1200} = \tilde{\Psi}_{1200} + (c + 6\psi_{3000})\zeta_2, \quad \Psi_{0300} = \tilde{\Psi}_{0300},$$

and

$$3c + 12\psi_{3000} = \langle 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0200}) + \mathbf{R}_{3,0}(\zeta_1, \zeta_1, \zeta_1) - 2\tilde{\Psi}_{1200}, \zeta_2^* \rangle.$$

From (D.17) and (D.18) we obtain

$$\Psi_{2010} = \tilde{\Psi}_{2010} + (3\psi_{3000} - 2c)\zeta_2 + \psi_{2010}\zeta_0, \quad \Psi_{1110} = \tilde{\Psi}_{1110} + 2\psi_{2010}\zeta_1,$$

and

$$\begin{aligned} 12\psi_{3000} = & \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0110}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{1010}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{1100}), \zeta_2^* \rangle \\ & + \langle 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \zeta_2), \zeta_2^* \rangle - \langle 2\tilde{\Psi}_{2010} + 2\tilde{\Psi}_{1200} + 2b\Psi_{0020}, \zeta_2^* \rangle. \end{aligned}$$

We conclude that

$$\begin{aligned}
3c = & \langle 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0200}) - 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{0110}) - 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{1010}), \zeta_2^* \rangle \\
& - \langle 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{1100}) - \mathbf{R}_{3,0}(\zeta_1, \zeta_1, \zeta_1) + 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \zeta_2), \zeta_2^* \rangle \\
& + \langle 2\tilde{\Psi}_{2010} + 2b\Psi_{0020}, \zeta_2^* \rangle.
\end{aligned} \tag{D.24}$$

Furthermore, from the equations (D.19) and (D.20) we can find

$$\Psi_{1020} = \tilde{\Psi}_{1020} + 2\psi_{2010}\zeta_2, \quad \Psi_{0210} = \tilde{\Psi}_{0210} + 2\psi_{2010}\zeta_2,$$

the solvability condition for (D.21) gives

$$4\psi_{2010} = \langle 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{0020}) + 2\mathbf{R}_{2,0}(\zeta_2, \Psi_{0110}) + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_2, \zeta_2), \zeta_2^* \rangle,$$

and from (D.21) and (D.22) we can also determine Ψ_{0120} and Ψ_{0030} .

D.2 $(i\omega)^2$ Normal Form in Infinite Dimensions

We show below how to compute the principal coefficients in the normal form (3.25) when starting from an infinite-dimensional system of the form (3.12). We assume that the parameter μ is real and that the system (3.12) possesses a reversibility symmetry \mathbf{S} and satisfies the hypotheses of Theorems 3.3 and 3.15 in Chapter 2. We further assume that the spectrum of the linear operator \mathbf{L} is such that $\sigma_0 = \{\pm i\omega\}$, where $\pm i\omega$ are algebraically double and geometrically simple eigenvalues. Then the four-dimensional reduced system satisfies the hypotheses of Lemma 3.17, so that its normal form is given by (3.25).

We proceed as in Section 3.4, and in the previous cases. In equality (4.3) in Chapter 3, we take $v_0 = A\zeta_0 + B\zeta_1 + \overline{A}\zeta_0 + \overline{B}\zeta_1$, and then write

$$u = A\zeta_0 + B\zeta_1 + \overline{A}\zeta_0 + \overline{B}\zeta_1 + \tilde{\Psi}(A, B, \overline{A}, \overline{B}, \mu) \tag{D.25}$$

where $\tilde{\Psi}$ takes values in \mathcal{X} . With the notations from Section 3.2.3, we consider the Taylor expansion (1.15) of \mathbf{R} , and the expansion of $\tilde{\Psi}$,

$$\tilde{\Psi}(A, B, \overline{A}, \overline{B}, \mu) = \sum_{1 \leq r+s+q+l+m \leq p} A^r B^s \overline{A}^q \overline{B}^l \mu^m \Psi_{rsqlm},$$

where

$$\Psi_{rsql0} = 0, \quad \text{for } r+s+q+l = 1.$$

Using the reversibility symmetry we find that

$$\tilde{\Psi}(\overline{A}, -\overline{B}, A, B, \mu) = \mathbf{S}\tilde{\Psi}(A, B, \overline{A}, \overline{B}, \mu),$$

and

$$\mathbf{S}\Psi_{rsqlm} = (-1)^{s+l}\Psi_{qlrsm}, \quad \Psi_{rsqlm} = \overline{\Psi}_{qlrsm}.$$

Identity (4.4) in Chapter 3, is in this case

$$\begin{aligned}
 & (i\omega A + B)\partial_A \Psi + i\omega B\partial_B \Psi + (-i\omega\bar{A} + \bar{B})\partial_{\bar{A}} \Psi - i\omega\bar{B}\partial_{\bar{B}} \Psi \\
 & + \left(iA(\zeta_0 + \partial_A \Psi) - i\bar{A}(\bar{\zeta}_0 + \partial_{\bar{A}} \Psi) \right) P \\
 & + (\zeta_1 + \partial_B \Psi)(iBP + AQ) + (\bar{\zeta}_1 + \partial_{\bar{B}} \Psi)(-i\bar{B}P + \bar{A}Q) \\
 & = \mathbf{L}\Psi + \mathbf{R}(A\zeta_0 + B\zeta_1 + \bar{A}\bar{\zeta}_0 + \bar{B}\bar{\zeta}_1 + \Psi, \mu).
 \end{aligned}$$

Using the expansions of \mathbf{R} , $\tilde{\Psi}$, P , and Q , we find at orders μ , $A\mu$, and $B\mu$, the equalities

$$0 = \mathbf{L}\Psi_{00001} + \mathbf{R}_{0,1}, \quad (\text{D.26})$$

$$a\zeta_1 + i\alpha\zeta_0 = (\mathbf{L} - i\omega)\Psi_{10001} + \mathbf{R}_{1,1}\zeta_0 + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{00001}), \quad (\text{D.27})$$

$$i\alpha\zeta_1 + \Psi_{10001} = (\mathbf{L} - i\omega)\Psi_{01001} + \mathbf{R}_{1,1}\zeta_1 + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{00001}), \quad (\text{D.28})$$

and at orders A^2 , $A\bar{A}$, AB , $A\bar{B}$, B^2 , $B\bar{B}$ we have:

$$0 = (\mathbf{L} - 2i\omega)\Psi_{20000} + \mathbf{R}_{2,0}(\zeta_0, \zeta_0), \quad (\text{D.29})$$

$$0 = \mathbf{L}\Psi_{10100} + 2\mathbf{R}_{2,0}(\zeta_0, \bar{\zeta}_0), \quad (\text{D.30})$$

$$2\Psi_{20000} = (\mathbf{L} - 2i\omega)\Psi_{11000} + 2\mathbf{R}_{2,0}(\zeta_0, \zeta_1), \quad (\text{D.31})$$

$$\Psi_{10100} = \mathbf{L}\Psi_{10010} + 2\mathbf{R}_{2,0}(\zeta_0, \bar{\zeta}_1), \quad (\text{D.32})$$

$$\Psi_{11000} = (\mathbf{L} - 2i\omega)\Psi_{02000} + \mathbf{R}_{2,0}(\zeta_1, \zeta_1), \quad (\text{D.33})$$

$$\Psi_{10010} + \Psi_{01100} = \mathbf{L}\Psi_{01010} + 2\mathbf{R}_{2,0}(\zeta_1, \bar{\zeta}_1). \quad (\text{D.34})$$

At orders A^3 , A^2B , AB^2 , and B^3 we obtain:

$$0 = (\mathbf{L} - 3i\omega)\Psi_{30000} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{20000}) + \mathbf{R}_{3,0}(\zeta_0, \zeta_0, \zeta_0), \quad (\text{D.35})$$

$$\begin{aligned}
 3\Psi_{30000} &= (\mathbf{L} - 3i\omega)\Psi_{21000} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{20000}) \\
 &+ 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \zeta_1),
 \end{aligned} \quad (\text{D.36})$$

$$\begin{aligned}
 2\Psi_{21000} &= (\mathbf{L} - 3i\omega)\Psi_{12000} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{02000}) \\
 &+ 3\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \zeta_1),
 \end{aligned} \quad (\text{D.37})$$

$$\Psi_{12000} = (\mathbf{L} - 3i\omega)\Psi_{03000} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{02000}) + \mathbf{R}_{3,0}(\zeta_1, \zeta_1, \zeta_1), \quad (\text{D.38})$$

and finally at orders $A^2\bar{A}$, $A^2\bar{B}$, $A\bar{A}B$, $\bar{A}B^2$, $AB\bar{B}$, and $B^2\bar{B}$ we find:

$$\begin{aligned}
 b\zeta_1 + i\beta\zeta_0 &= (\mathbf{L} - i\omega)\Psi_{20100} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{10100}) \\
 &+ 2\mathbf{R}_{2,0}(\bar{\zeta}_0, \Psi_{20000}) + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \bar{\zeta}_0),
 \end{aligned} \quad (\text{D.39})$$

$$\begin{aligned}
 \frac{ic}{2}\zeta_1 - \frac{\gamma}{2}\zeta_0 + \Psi_{20100} &= (\mathbf{L} - i\omega)\Psi_{20010} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{10010}) \\
 &+ 2\mathbf{R}_{2,0}(\bar{\zeta}_1, \Psi_{20000}) + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \bar{\zeta}_1),
 \end{aligned} \quad (\text{D.40})$$

$$\begin{aligned} \left(i\beta - \frac{ic}{2}\right) \zeta_1 + \frac{\gamma}{2} \zeta_0 + 2\Psi_{20100} &= (\mathbf{L} - i\omega)\Psi_{11100} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{01100}) \\ &\quad + 2\mathbf{R}_{2,0}(\bar{\zeta}_0, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{10100}) \\ &\quad + 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \bar{\zeta}_0), \end{aligned} \quad (\text{D.41})$$

$$\begin{aligned} \frac{\gamma}{2} \zeta_1 + \Psi_{11100} &= (\mathbf{L} - i\omega)\Psi_{02100} + 2\mathbf{R}_{2,0}(\bar{\zeta}_0, \Psi_{02000}) \\ &\quad + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{01100}) + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \bar{\zeta}_0), \end{aligned} \quad (\text{D.42})$$

$$\begin{aligned} -\frac{\gamma}{2} \zeta_1 + 2\Psi_{20010} + \Psi_{11100} &= (\mathbf{L} - i\omega)\Psi_{11010} + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{01010}) \\ &\quad + 2\mathbf{R}_{2,0}(\bar{\zeta}_1, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{10010}) \\ &\quad + 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \bar{\zeta}_1), \end{aligned} \quad (\text{D.43})$$

$$\begin{aligned} \Psi_{11010} + \Psi_{02100} &= (\mathbf{L} - i\omega)\Psi_{02010} + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{01010}) \\ &\quad + 2\mathbf{R}_{2,0}(\bar{\zeta}_1, \Psi_{02000}) + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \bar{\zeta}_1). \end{aligned} \quad (\text{D.44})$$

Notice that due to the symmetry properties of $\mathbf{R}_{p,q}$, ζ_0 , and ζ_1 we have

$$\mathbf{S}\mathbf{R}_{0,1} = -\mathbf{R}_{0,1}, \quad \mathbf{S}\mathbf{R}_{1,1}\zeta_0 = -\overline{\mathbf{R}_{1,1}\zeta_0}, \quad \mathbf{S}\mathbf{R}_{1,1}\zeta_1 = \overline{\mathbf{R}_{1,1}\zeta_1},$$

which, together with the symmetry properties for Ψ_{qlrsm} , imply that applying the symmetry \mathbf{S} to both sides of (D.27), (D.39), (D.42), and (D.43), we obtain the opposite of the complex conjugate of these equalities, whereas by applying it to (D.28), (D.40), (D.41), and (D.44), we find the complex conjugate of these equalities.

First, notice that the invertibility of the operators \mathbf{L} , $(\mathbf{L} - 2i\omega)$, and $(\mathbf{L} - 3i\omega)$, lets us solve the equations (D.26), (D.29)–(D.34), and (D.35)–(D.38), and determine Ψ_{00001} , Ψ_{20000} , Ψ_{10100} , Ψ_{11000} , Ψ_{10010} , Ψ_{02000} , Ψ_{01010} , Ψ_{30000} , Ψ_{21000} , Ψ_{12000} , and Ψ_{03000} .

Next, consider the vector ζ_1^* orthogonal to the range of $\mathbf{L} - i\omega$, constructed as in the other cases, such that

$$\langle \zeta_0, \zeta_1^* \rangle = 0, \quad \langle \bar{\zeta}_0, \zeta_1^* \rangle = 0, \quad \langle \zeta_1, \zeta_1^* \rangle = 1, \quad \langle \bar{\zeta}_1, \zeta_1^* \rangle = 0,$$

and

$$\mathbf{S}^* \zeta_1^* = -\bar{\zeta}_1^*.$$

Then from the equations (D.27) and (D.28) we find

$$a = \langle \mathbf{R}_{1,1}\zeta_0 + 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{00001}), \zeta_1^* \rangle \quad (\text{D.45})$$

and

$$\begin{aligned} \Psi_{10001} &= \tilde{\Psi}_{10001} + i\alpha\zeta_1, \\ 2i\alpha &= \langle \mathbf{R}_{1,1}\zeta_1 + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{00001}) - \tilde{\Psi}_{10001}, \zeta_1^* \rangle. \end{aligned} \quad (\text{D.46})$$

Taking into account the fact that $\langle \mathbf{S}u, \mathbf{S}^*v \rangle = \langle u, v \rangle$ for any $u \in \mathcal{X}$ and $v \in \mathcal{X}^*$, it is not difficult to check that a and α are real numbers.

Next, equation (D.39) gives

$$b = \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{10100}) + 2\mathbf{R}_{2,0}(\overline{\zeta_0}, \Psi_{20000}) + 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \overline{\zeta_0}), \zeta_1^* \rangle, \quad (\text{D.47})$$

and

$$\Psi_{20100} = \tilde{\Psi}_{20100} + i\beta \zeta_1 + \psi_{20100} \zeta_0,$$

with a constant $\psi_{20100} \in \mathbb{R}$, which will be determined later. The equations (D.40) and (D.41) now give

$$\begin{aligned} i\beta + \frac{ic}{2} &= \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{10010}) + 2\mathbf{R}_{2,0}(\overline{\zeta_1}, \Psi_{20000}), \zeta_1^* \rangle \\ &\quad + \langle 3\mathbf{R}_{3,0}(\zeta_0, \zeta_0, \overline{\zeta_1}) - \tilde{\Psi}_{20100}, \zeta_1^* \rangle, \\ 3i\beta - \frac{ic}{2} &= \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{01100}) + 2\mathbf{R}_{2,0}(\overline{\zeta_0}, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{10100}), \zeta_1^* \rangle \\ &\quad + \langle 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \overline{\zeta_0}) - 2\tilde{\Psi}_{20100}, \zeta_1^* \rangle, \end{aligned}$$

which determine the real coefficients β and c , and

$$\begin{aligned} \Psi_{20010} &= \tilde{\Psi}_{20010} + \left(\psi_{20100} - \frac{\gamma}{2} \right) \zeta_1, \\ \Psi_{11100} &= \tilde{\Psi}_{11100} + \left(2\psi_{20100} + \frac{\gamma}{2} \right) \zeta_1 + \psi_{11100} \zeta_0, \end{aligned}$$

with another constant ψ_{11100} determined later.

Finally, the equations (D.42)–(D.44) let us determine γ , ψ_{20100} , and ψ_{11100} . Indeed, (D.42) gives

$$\begin{aligned} \gamma + 2\psi_{20100} &= \langle 2\mathbf{R}_{2,0}(\overline{\zeta_0}, \Psi_{02000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{01100}), \zeta_1^* \rangle \\ &\quad + \langle 3\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \overline{\zeta_0}) - \tilde{\Psi}_{11100}, \zeta_1^* \rangle, \end{aligned}$$

and

$$\Psi_{02100} = \tilde{\Psi}_{02100} + \psi_{11100} \zeta_1,$$

whereas (D.43) leads to

$$\begin{aligned} 4\psi_{20100} - \gamma &= \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{01010}) + 2\mathbf{R}_{2,0}(\overline{\zeta_1}, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{10010}), \zeta_1^* \rangle \\ &\quad + \langle 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \overline{\zeta_1}) - 2\tilde{\Psi}_{20010} - \tilde{\Psi}_{11100}, \zeta_1^* \rangle, \end{aligned}$$

and

$$\Psi_{11010} = \tilde{\Psi}_{11010} + \psi_{11100} \zeta_1.$$

This determines γ and ψ_{20100} . In particular, we obtain

$$\begin{aligned} 3\gamma &= \langle 4\mathbf{R}_{2,0}(\overline{\zeta_0}, \Psi_{02000}) + 4\mathbf{R}_{2,0}(\zeta_1, \Psi_{01100}) + 6\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \overline{\zeta_0}), \zeta_1^* \rangle \\ &\quad - \langle 2\mathbf{R}_{2,0}(\zeta_0, \Psi_{01010}) + 2\mathbf{R}_{2,0}(\overline{\zeta_1}, \Psi_{11000}) + 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{10010}), \zeta_1^* \rangle \\ &\quad - \langle 6\mathbf{R}_{3,0}(\zeta_0, \zeta_1, \overline{\zeta_1}) - 2\tilde{\Psi}_{20010} + \tilde{\Psi}_{11100}, \zeta_1^* \rangle. \end{aligned}$$

Finally the equation (D.44) gives

$$2\psi_{11100} = \langle 2\mathbf{R}_{2,0}(\zeta_1, \Psi_{01010}) + 2\mathbf{R}_{2,0}(\overline{\zeta_1}, \Psi_{02000}) + 3\mathbf{R}_{3,0}(\zeta_1, \zeta_1, \overline{\zeta_1}), \zeta_1^* \rangle \\ - \langle \tilde{\Psi}_{11010} + \tilde{\Psi}_{02100}, \zeta_1^* \rangle,$$

and completes the computation of the normal form.

References

1. R.A. Adams. *Sobolev spaces*. Pure and Applied Mathematics, Vol. 65. Academic Press, New York-London, 1975.
2. S. Agmon. *Lectures on elliptic boundary value problems*. Van Nostrand Mathematical Studies, No. 2, D. Van Nostrand Co., Inc., Princeton, NJ, Toronto-London, 1965.
3. A. Andronov, E. Leontovich. Some cases of dependence of limit cycles on a parameter. (Russian) *J. State Univ. Gorki* 6 (1937), 3–24.
4. V.I. Arnold. *Geometrical methods in the theory of ordinary differential equations*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], 250. Springer-Verlag, New York-Berlin, 1983.
5. M. Barrandon. Résonance 1 : 2 pour les champs de vecteurs réversibles. *C. R. Math. Acad. Sci. Paris* 334 (2002), 1, 7–10.
6. J.T. Beale. Exact solitary water waves with capillary ripples at infinity. *Comm. Pure Appl. Math.* 44 (1991), 2, 211–257.
7. G.R. Belickii. Normal forms in relation to the filtering action of a group. (Russian) *Trudy Moskov. Mat. Obshch.* 40 (1979), 3–46.
8. D. Bensimon, P. Kolodner, C.M. Surko, H. Williams, V. Croquette. Competing and coexisting dynamical states of travelling wave convection in an annulus. *J. Fluid Mech.* 217 (1990), 441–467.
9. D. Brézis. Perturbations singulières et problèmes d'évolution avec défaut d'ajustement. *C. R. Acad. Sci. Paris Sér. A-B* 276 (1973), A1597–A1600.
10. B. Buffoni, J. Toland. *Analytic theory of global bifurcation. An introduction*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2003.
11. J.M. Burgers. A mathematical model illustrating the theory of turbulence. *Adv. Appl. Mech.* 1 (1948), 171–199.
12. J. Carr. *Applications of centre manifold theory*. Applied Mathematical Sciences, 35. Springer-Verlag, New York-Berlin, 1981.
13. A.R. Champneys. Homoclinic orbits in reversible systems and their applications in mechanics, fluids and optics. Time-reversal symmetry in dynamical systems (Coventry, 1996). *Phys. D* 112 (1998), 1-2, 158–186.
14. S. Chandrasekhar. *Hydrodynamic and hydromagnetic stability*. The International Series of Monographs on Physics. Clarendon Press, Oxford, 1961.
15. P. Chossat, G. Iooss. *The Couette–Taylor problem*. Applied Mathematical Sciences, 102. Springer-Verlag, New York, 1994.
16. P. Chossat, R. Lauterbach. *Methods in equivariant bifurcations and dynamical systems*. Advanced Series in Nonlinear Dynamics, 15. World Scientific Publishing Co., Inc., River Edge, NJ, 2000.
17. S.N. Chow, J. K. Hale. *Methods of bifurcation theory*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], 251. Springer-Verlag, New York-Berlin, 1982.

18. P. Collet, J.P. Eckmann. *Instabilities and fronts in extended systems*. Princeton Series in Physics. Princeton University Press, Princeton, NJ, 1990.
19. P. Coullet, G. Iooss. Instabilities of one-dimensional cellular patterns. *Phys. Rev. Lett.* 64 (1990), 8, 866–869.
20. R. Cushman, J.A. Sanders. Splitting algorithm for nilpotent normal forms. *Dynam. Stability Systems* 2 (1987), 3–4, 235–246.
21. G. Da Prato, P. Grisvard. Sommes d'opérateurs linéaires et équations différentielles opérationnelles. *J. Math. Pures Appl.* (9) 54 (1975), 3, 305–387.
22. F. Dias, G. Iooss. Capillary-gravity solitary waves with damped oscillations. *Phys. D* 65 (1993), 4, 399–423.
23. F. Dias, G. Iooss. Water-waves as a spatial dynamical system. *Handbook of mathematical fluid dynamics*, Vol. II 443–499, North-Holland, Amsterdam, 2003.
24. C. Elphick, G. Iooss, E. Tirapegui. Normal form reduction for time-periodically driven differential equations. *Phys. Lett. A* 120 (1987), 9, 459–463.
25. C. Elphick, E. Tirapegui, M.E. Brachet, P. Coullet, G. Iooss. A simple global characterization for normal forms of singular vector fields. *Phys. D* 29 (1987), 1–2, 95–127.
26. J.-M. Gambaudo. Perturbation of a Hopf bifurcation by an external time-periodic forcing. *J. Differential Equations* 57 (1985), 2, 172–199.
27. S.A. van Gils, M. Krupa, W.F. Langford. Hopf bifurcation with nonsemisimple 1 : 1 resonance. *Nonlinearity* 3 (1990), 3, 825–850.
28. L. Yu. Glebsky, L.M. Lerman. On small stationary localized solutions for the generalized 1-D Swift-Hohenberg equation. *Chaos* 5 (1995), 2, 424–431.
29. M. Golubitsky, D.G. Schaeffer. *Singularities and groups in bifurcation theory*. Vol. I. Applied Mathematical Sciences, 51. Springer-Verlag, New York, 1985.
30. M. Golubitsky, I. Stewart, D.G. Schaeffer. *Singularities and groups in bifurcation theory*. Vol. II. Applied Mathematical Sciences, 69. Springer-Verlag, New York, 1988.
31. R. Grimshaw, G. Iooss. Solitary waves of a coupled Korteweg-de Vries system. Nonlinear waves: computation and theory, II (Athens, GA, 2001). *Math. Comput. Simulation* 62 (2003), 1–2, 31–40.
32. D.M. Grobman. Homeomorphism of systems of differential equations. (Russian) *Dokl. Akad. Nauk SSSR* 128 (1959), 880–881.
33. M.D. Groves. Three-dimensional travelling gravity-capillary water waves. *GAMM-Mitt.* 30 (2007), 1, 8–43.
34. M.D. Groves, M. Haragus. A bifurcation theory for three-dimensional oblique travelling gravity-capillary water waves. *J. Nonlinear Sci.* 13 (2003), 4, 397–447.
35. M.D. Groves, A. Mielke. A spatial dynamics approach to three-dimensional gravity-capillary steady water waves. *Proc. Roy. Soc. Edinburgh Sect. A* 131 (2001), 1, 83–136.
36. M.D. Groves, B. Sandstede. A plethora of three-dimensional periodic travelling gravity-capillary water waves with multipulse transverse profiles. *J. Nonlinear Sci.* 14 (2004), 3, 297–340.
37. M.D. Groves, S.M. Sun. Fully localised solitary-wave solutions of the three-dimensional gravity-capillary water-wave problem. *Arch. Ration. Mech. Anal.* 188 (2008), 1, 1–91.
38. J. Guckenheimer, P. Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*. Applied Mathematical Sciences, 42. Springer-Verlag, New York, 1983.
39. J.K. Hale, H. Koçak. *Dynamics and bifurcations*. Texts in Applied Mathematics, 3. Springer-Verlag, New York, 1991.
40. M. Haragus, T. Kapitula. Spots and stripes in NLS-type equations with nearly one-dimensional potentials. *Math. Methods Appl. Sci.* (to appear).
41. M. Haragus, K. Kirchgässner. Three-dimensional steady capillary-gravity waves. *Ergodic theory, analysis, and efficient simulation of dynamical systems*, 363–397, Springer, Berlin, 2001.
42. M. Haragus, A. Scheel. Corner defects in almost planar interface propagation. *Ann. Inst. H. Poincaré, Anal. Non Linéaire* 23 (2006), 3, 283–329.
43. M. Haragus, A. Scheel. Almost planar waves in anisotropic media. *Comm. Partial Differential Equations* 31 (2006), 4–6, 791–815.

44. M. Haragus, A. Scheel. A bifurcation approach to non-planar traveling waves in reaction-diffusion systems. *GAMM-Mitt.* 30 (2007), 1, 75–95.
45. M. Haragus, A. Scheel. Interfaces between rolls in the Swift-Hohenberg equation. *Int. J. Dyn. Syst. Differ. Equ.* 1 (2007), 2, 89–97.
46. P. Hartman. A lemma in the theory of structural stability of differential equations. *Proc. Amer. Math. Soc.* 11 (1960), 610–620.
47. D. Henry. *Geometric theory of semilinear parabolic equations*. Lecture Notes in Mathematics, 840. Springer-Verlag, Berlin-New York, 1981.
48. M.W. Hirsch, C.C. Pugh. Stable manifolds and hyperbolic sets. Global Analysis. *Proc. Symp. Pure Math., Vol. XIV, Berkeley, CA* (1968) pp. 133–163; Amer. Math. Soc., Providence, R.I., 1970.
49. A.J. Homburg, B. Sandstede. Homoclinic and heteroclinic bifurcations in vector fields. *Handbook of Dynamical Systems*, Vol. III (to appear).
50. E. Hopf. Abzweigung einer periodischen Lösung von einer stationären Lösung eines Differentialsystems. *Ber. Verh. Sächs. Akad. Wiss. Leipzig. Math.-Nat. Kl.* 95 (1943), 1, 3–22.
51. K. Ikeda, K. Murota. *Imperfect Bifurcation in Structures and Materials*. Applied Math. Sci., 149, Springer, New York, 2002.
52. G. Iooss. Théorie non linéaire de la stabilité des écoulements laminaires dans le cas de “l’échange des stabilités”. *Arch. Rational Mech. Anal.* 40 (1970/1971), 166–208.
53. G. Iooss. Global characterization of the normal form for a vector field near a closed orbit. *J. Diff. Equ.* 76 (1988), 1, 47–76.
54. G. Iooss. A codimension 2 bifurcation for reversible vector fields. Normal forms and homoclinic chaos (Waterloo, ON, 1992), *Fields Inst. Com.* 4 (1995), 201–217.
55. G. Iooss. Travelling waves in the Fermi–Pasta–Ulam lattice. *Nonlinearity* 13 (2000), 3, 849–866.
56. G. Iooss, M. Adelmeyer. *Topics in bifurcation theory and applications*. Second ed. Advanced Series in Nonlinear Dynamics, 3. World Scientific Publishing, River Edge, NJ, 1998.
57. G. Iooss, G. James. Localized waves in nonlinear oscillator chains. *Chaos* 15 (2005), 1, 015113, 15 pp.
58. G. Iooss, D.D. Joseph. *Elementary stability and bifurcation theory*. Second ed. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1990.
59. G. Iooss, K. Kirchgässner. Water waves for small surface tension: an approach via normal form. *Proc. Roy. Soc. Edinburgh A* 122 (1992), 3–4, 267–299.
60. G. Iooss, K. Kirchgässner. Travelling waves in a chain of coupled nonlinear oscillators. *Comm. Math. Phys.* 211 (2000), 2, 439–464.
61. G. Iooss, E. Lombardi. Normal forms with exponentially small remainder: application to homoclinic connections for the reversible $0^{2+}i\omega$ resonance. *C. R. Math. Acad. Sci. Paris* 339 (2004), 12, 831–838.
62. G. Iooss, E. Lombardi. Polynomial normal forms with exponentially small remainder for analytic vector fields. *J. Diff. Equ.* 212 (2005), 1, 1–61.
63. G. Iooss, E. Lombardi. Approximated invariant manifold up to exponentially small terms. *J. Diff. Equ.* 248 (2010), 6, 1410–1431.
64. G. Iooss, J. Los. Bifurcation of spatially quasi-periodic solutions in hydrodynamic stability problems. *Nonlinearity* 3 (1990), 851–871.
65. G. Iooss, A. Mielke. Bifurcating time-periodic solutions of Navier-Stokes equations in infinite cylinders. *J. Nonlinear Sci.* 1 (1991), 107–146.
66. G. Iooss, A. Mielke. Time-periodic Ginzburg-Landau equations for one-dimensional patterns with large wave length. *Z. Angew. Math. Phys.* 43 (1992), 1, 125–138.
67. G. Iooss, A. Mielke, Y. Demay. Theory of steady Ginzburg-Landau equation, in hydrodynamic stability problems. *European J. Mech. B Fluids* 8 (1989), 3, 229–268.
68. G. Iooss, M.-C. Pérouème. Perturbed homoclinic solutions in reversible 1 : 1 resonance vector fields. *J. Diff. Equ.* 102 (1993), 1, 62–88.
69. G. Iooss, P. Plotnikov. Small divisor problem in the theory of three-dimensional water gravity waves. *Mem. Amer. Math. Soc.* 200 (2009), 940, 1–128.

70. G. Iooss, M. Rossi. Hopf bifurcation in the presence of spherical symmetry: analytical results. *SIAM J. Math. Anal.* 20 (1989), 3, 511–532.
71. C.G.J. Jacobi. Über die Figur des Gleichgewichts. *Ann. Phys. Chem.* 33 (1834) 229–233.
72. G. James. Centre manifold reduction for quasilinear discrete systems. *J. Nonlinear Sci.* 13 (2003), 1, 27–63.
73. G. James. Internal travelling waves in the limit of a discontinuously stratified fluid. *Arch. Ration. Mech. Anal.* 160 (2001), 1, 41–90.
74. A. Joets. Apollonios, premier géomètre des singularités. *Quadrature* 66 (2007), 37–41
75. D.D. Joseph. *Stability of fluid motions*. I and II. Springer Tracts in Natural Philosophy, Vol. 27 and 28. Springer-Verlag, Berlin-New York, 1976.
76. T. Kato. *Perturbation theory for linear operators*. Reprint of the 1980 ed. Classics in Mathematics. Springer-Verlag, Berlin, 1995.
77. A. Kelley. The stable, center-stable, center, center-unstable, unstable manifolds. *J. Differential Equations* 3 (1967), 546–570.
78. H. Kielhöfer, K. Kirchgässner. Stability and bifurcation in fluid dynamics. Rocky Mountain Consortium Symposium on Nonlinear Eigenvalue Problems (Santa Fe, N.M., 1971). *Rocky Mountain J. Math.* 3 (1973), 275–318.
79. H. Kielhöfer. *Bifurcation theory. An introduction with applications to PDEs*. Applied Mathematical Sciences, 156. Springer-Verlag, New York, 2004.
80. K. Kirchgässner. Wave-solutions of reversible systems and applications. *J. Diff. Equ.* 45 (1982), 1, 113–127.
81. K. Kirchgässner. Nonlinearly resonant surface waves and homoclinic bifurcation. *Adv. Appl. Mech.* 26 (1988), 135–181.
82. P. Kirrmann. Reduktion nichtlinearer elliptischer Systeme in Zylindergebieten unter Verwendung von optimaler Regularität in Hölder Räumen. PhD thesis, Universität Stuttgart, 1991.
83. E.L. Koschmieder. *Bénard cells and Taylor vortices*. Cambridge Monographs on Mechanics and Applied Mathematics. Cambridge University Press, New York, 1993.
84. Y.A. Kuznetsov. *Elements of applied bifurcation theory*. Third ed. Applied Mathematical Sciences, 112. Springer-Verlag, New York, 2004.
85. O.A. Ladyzhenskaya. *The mathematical theory of viscous incompressible flow*. Revised English ed. Translated from the Russian, Gordon and Breach Science Pub., New York-London 1963.
86. L. Landau. On the problem of turbulence. *C. R. (Doklady) Acad. Sci. URSS (N.S.)* 44 (1944), 311–314.
87. O.E. Lanford. Bifurcation of periodic solutions into invariant tori: the work of Ruelle and Takens. *Nonlinear problems in the physical sciences and biology*. Proceedings of a Battelle Summer Institute, Seattle, Wash., July 3–28, 1972. I. Stakgold, D. D. Joseph and D. H. Sattinger eds. Lecture Notes in Mathematics, Vol. 322. Springer-Verlag, Berlin-New York, 1973.
88. W.F. Langford, G. Iooss. Interactions of Hopf and pitchfork bifurcations. Bifurcation problems and their numerical solution (Proc. Workshop, Univ. Dortmund, Dortmund, 1980), *Internat. Ser. Numer. Math.* 54 (1980), 103–134.
89. V.G. LeBlanc, W.F. Langford. Classification and unfoldings of 1 : 2 resonant Hopf bifurcation. *Arch. Rational Mech. Anal.* 136 (1996), 4, 305–357.
90. T. Levi-Civita. Détermination rigoureuse des ondes permanentes d’amplitude finie. *Math. Ann.* 93 (1925), 1, 264–314.
91. E. Lombardi. Orbits homoclinic to exponentially small periodic orbits for a class of reversible systems. Application to water waves. *Arch. Rational Mech. Anal.* 137 (1997), 3, 227–304.
92. E. Lombardi. Non-persistence of homoclinic connections for perturbed integrable reversible systems. *J. Dyn. Diff. Equ.* 11 (1999), 1, 129–208.
93. E. Lombardi. *Oscillatory integrals and phenomena beyond all algebraic orders. With applications to homoclinic orbits in reversible systems*. Lecture Notes in Mathematics, 1741. Springer-Verlag, Berlin, 2000.
94. J.E. Marsden, M. McCracken. *The Hopf bifurcation and its applications*. Applied Mathematical Sciences, Vol. 19. Springer-Verlag, New York, 1976.

95. A. Mielke. A reduction principle for nonautonomous systems in infinite-dimensional spaces. *J. Diff. Equ.* 65 (1986), 1, 68–88.
96. A. Mielke. Über maximale L^p -Regularität für Differentialgleichungen in Banach- und Hilbert-Räumen. *Math. Ann.* 277 (1987), 1, 121–133.
97. A. Mielke. Reduction of quasilinear elliptic equations in cylindrical domains with applications. *Math. Methods Appl. Sci.* 10 (1988), 1, 51–66.
98. A. Mielke. On nonlinear problems of mixed type: a qualitative theory using infinite-dimensional center manifolds. *J. Dynam. Diff. Equ.* 4 (1992), 3, 419–443.
99. L.A. Ostrowsky. Nonlinear internal waves in a rotating ocean. *Okeanologia* 18 (1978), 181–191.
100. R. L. Pego, J. R. Quintero. A host of traveling waves in a model of three-dimensional water-wave dynamics. *J. Nonlinear Sci.* 12 (2002), 1, 59–83.
101. V. Pliss. A reduction principle in the theory of stability of motion. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 28 (1964) 1297–1324.
102. H. Poincaré. *Les méthodes nouvelles de la mécanique céleste*. Tome I. Reprint of the 1892 original. Les Grands Classiques Gauthier-Villars. Librairie Scientifique et Technique Albert Blanchard, Paris, 1987.
103. H. Poincaré. Sur l'équilibre d'une masse fluide animée d'un mouvement de rotation. *Acta Math.* 7 (1885), 1, 259–380.
104. B. van der Pol. Forced oscillations in a circuit with nonlinear resistance. *Phil. Mag.* 3 (1927), 65–80.
105. Y. Pomeau, A. Ramani, B. Grammaticos. Structural stability of the Korteweg-de Vries solitons under a singular perturbation. *Phys. D* 31 (1988), 1, 127–134.
106. I. Prigogine, R. Lefever. Symmetry breaking instabilities in dissipative systems. 2. *J. Chem. Phys.* 48 (1968), 4, 1695–1700.
107. C. Rorres. Completing Book II of Archimedes's on floating bodies. *Math. Intelligencer* 26 (2004), 3, 32–42.
108. D. Ruelle, F. Takens. On the nature of turbulence. *Comm. Math. Phys.* 20 (1971), 167–192, and 23 (1971), 343–344.
109. D.H. Sattinger. *Group-theoretic methods in bifurcation theory*. Lecture Notes in Mathematics, 762. Springer, Berlin, 1979.
110. S.H. Strogatz. *Nonlinear dynamics and chaos: With applications to physics, biology, chemistry, and engineering*. Studies in Nonlinearity, Perseus Books, Cambridge, 1994.
111. M.B. Sevryuk. *Reversible systems*. Lecture Notes in Mathematics, 1211. Springer-Verlag, Berlin, 1986.
112. J. Sijbrand. Studies in nonlinear stability and bifurcation theory. PhD Thesis, University of Utrecht, 1981.
113. V.S. Sorokin. On steady motions in a fluid heated from below. (Russian) *Akad. Nauk SSSR. Prikl. Mat. Meh.* 18 (1954), 197–204.
114. S.J. van Strien. Center manifolds are not C^∞ . *Math. Z.* 166 (1979), 2, 143–145.
115. S.M. Sun. Existence of a generalized solitary wave solution for water with positive Bond number less than $1/3$. *J. Math. Anal. Appl.* 156 (1991), 2, 471–504.
116. S.M. Sun, M.C. Shen. Exponentially small estimate for the amplitude of capillary ripples of a generalized solitary wave. *J. Math. Anal. Appl.* 172 (1993), 2, 533–566.
117. R. Tagg. The Couette–Taylor problem. *Nonlinear Science Today* 4 (1994), 2–25; see also: Couette–Taylor Reference Base, 1999, <http://carbon.cudenver.edu/rtagg/Special/SpecialWebMaterials.html>.
118. R. Temam. *Navier–Stokes equations. Theory and numerical analysis*. Studies in Mathematics and its Applications, Vol. 2. North-Holland Publishing Co., Amsterdam–New York–Oxford, 1977.
119. M.R. Ukhovskii, V.I. Yudovich. On the equations of steady-state convection. *Prikl. Mat. Meh.* 27, 295–300 (Russian); translated as *J. Appl. Math. Mech.* 27 (1963), 432–440.
120. A. Vanderbauwhede. Centre manifolds, normal forms and elementary bifurcations. *Dynamics reported*, Vol. 2, 89–169, Dynam. Report. Ser. Dynam. Systems Appl., 2, Wiley, Chichester, 1989.

121. A. Vanderbauwhede, S.A. van Gils. Center manifolds and contractions on a scale of Banach spaces. *J. Funct. Anal.* 72 (1987), 2, 209–224.
122. A. Vanderbauwhede, G. Iooss. Center manifold theory in infinite dimensions. *Dynamics reported: expositions in dynamical systems*, 125–163, Dynam. Report. Expositions Dynam. Systems (N.S.), 1, Springer, Berlin, 1992.
123. W. Velte. Stabilitätsverhalten und Verzweigung stationärer Lösungen der Navier-Stokesschen Gleichungen. *Arch. Rational Mech. Anal.* 16 (1964), 97–125.
124. L.-J. Wang. Homoclinic and heteroclinic orbits for the 0^2 or $0^2i\omega$ singularity in the presence of two reversibility symmetries. *Quart. Appl. Math.* 67 (2009), 1–38.
125. L.-J. Wang. Studies on 0^2 or $0^2i\omega$ singularity in the presence of two reversibility symmetries and on dynamics of the Swift–Hohenberg equation with Dirichlet–Neumann boundary conditions. PhD thesis, Université de Toulouse, 2008.
126. V.I. Yudovich. Secondary flows and fluid instability between rotating cylinders. *Prikl. Mat. Meh.* 30, 688–698 (Russian); translated as *J. Appl. Math. Mech.* 30 (1966), 822–833.
127. V.I. Yudovich. On the origin of convection. *Prikl. Mat. Meh.* 30, 1000–1005 (Russian); translated as *J. Appl. Math. Mech.* 30 (1966), 1193–1199.
128. V.I. Yudovich. Free convection and bifurcation. *Prikl. Mat. Meh.* 31, 101–111 (Russian); translated as *J. Appl. Math. Mech.* 31 (1967), 103–114.
129. V.I. Yudovich. *The linearization method in hydrodynamical stability theory*. Translations of Mathematical Monographs, 74. American Mathematical Society, Providence, RI, 1989.

Index

- $D(\mathbf{L})$, 280
- $H^m(\Omega)$, 285
- $H_0^m(\Omega)$, 286
- $H_{\text{per}}^m(0, \ell)$, 286
- $L^2(\Omega)$, 285
- $L_{\text{loc}}^2(\mathbb{R})$, 286
- $L_{\text{per}}^2(0, \ell)$, 286
- \mathbb{N}^* , 43
- \mathbf{L}^* , 283
- $\mathcal{C}(\mathcal{X})$, 280
- $\mathcal{L}(\mathcal{X})$, 279
- $\mathcal{L}(\mathcal{Z}, \mathcal{X})$, 279
- $G(\mathbf{L})$, 279
- $\text{im } \mathbf{L}$, 279
- $\ker \mathbf{L}$, 279
- $\rho(\mathbf{L})$, 280
- $\sigma(\mathbf{L})$, 280
- $\sigma_\infty(\mathbf{L})$, 281
- h.o.t.*, 70
- $\text{ind}(\mathbf{L})$, 284
- Exercise*
 - Chapter 2, 33, 49, 59, 70, 77–79
 - Chapter 3, 100, 107, 111, 121, 132, 133, 142, 151, 155
 - Chapter 4, 165, 167, 169, 170, 178, 191, 192, 206, 232, 236
- bifurcation, 1
 - $(i\omega_1)(i\omega_2)$, 139
 - dynamic, 10
 - fold, 3
 - Hopf, 9, 14–16, 19, 22, 24, 84, 85, 120, 121, 124, 126, 127, 129, 147, 198, 214, 247, 248
 - parameter, 1
 - pitchfork, 5–8, 13, 27, 67, 69, 247, 253, 258
 - reversible, 157
 - reversible $(i\omega)^2$, 214, 223, 263
 - reversible $(i\omega_1)(i\omega_2)$, 226
 - reversible $(i\omega_1)^2(i\omega_2)$, 269
 - reversible $0(i\omega)$, 195
 - reversible 00^2 , 193
 - reversible 0^20^2 , 234
 - reversible 0^{2+} , 160, 164, 208, 263, 278
 - reversible $0^{2+}(i\omega)$, 198, 204, 209, 263
 - reversible 0^{2-} , 172, 177
 - reversible $0^{2-}(i\omega)$, 210
 - reversible 0^{3+} , 180, 190
 - reversible 0^{3-} , 191
 - reversible 0^{4+} , 229, 263
 - saddle-node, 2, 3, 5, 231, 274
 - steady, 10, 24, 27, 28, 73, 75, 246, 252
 - subcritical, 6
 - supercritical, 6, 13, 22, 69, 121
 - symmetry-breaking, 248
 - Takens–Bogdanov, 134, 160, 164, 172, 177
 - turning point, 3
- Bose–Einstein condensates, 207
- center manifold, 29
 - computation, 35, 39, 61, 122
 - continuous symmetry, 55, 57
 - equivariant systems, 53
 - global, 35
 - hypotheses, 31, 33, 36, 46, 51, 53–55, 58
 - local, 34
 - nonautonomous, 51, 55
 - parameter-dependent, 46
 - reversible systems, 54, 55
 - theorem, 34, 37, 46, 51, 53, 54, 57, 59
- codimension, 1
- eigenspace, 280
- eigenvalue, 280

- geometrically simple, 281
- semisimple, 281
- simple, 281
- eigenvector, 281
- equation
 - advance-delay differential, 276
 - Korteweg–de Vries, 233, 236
 - Kuramoto–Sivashinsky, 274
 - Landau, 247
 - parabolic, 41
 - parameter-dependent, 46
 - quasilinear, 36
 - reduced, 34, 49
 - Schrödinger, 206
 - semilinear, 36
 - Swift–Hohenberg, 70, 223
- FPU, 275
- Fredholm index, 284
- graph, 279
 - norm, 280
- KAM, 213
- KdV, 233, 236
- KG, 275
- lattice, 275
 - Fermi–Pasta–Ulam, 275
 - Klein–Gordon, 275
- locally Lipschitz boundary, 286
- Lyapunov–Schmidt method, 257
- multiplicity
 - algebraic, 281
 - geometric, 281
- NLS, 206
- normal form, 93
 - $(i\omega_1)(i\omega_2)$ bifurcation, 140
 - analytic vector fields, 152, 153
 - characterization, 94, 110
 - computation, 16, 20, 85, 102, 112, 114, 120, 122, 125, 135, 140, 149, 166, 177, 205, 225, 310, 315
 - equivariant systems, 117
 - Hopf bifurcation, 10, 16, 20, 85, 120, 124, 130, 147
 - hypotheses, 93, 109, 117, 118, 144
 - linear, 111
 - parameter-dependent, 109, 110
 - particular cases, 99, 100, 104–106, 108, 109
 - reversible 0^{2-} bifurcation, 177
 - reversible $(i\omega)^2$ bifurcation, 215, 315
 - reversible $(i\omega_1)(i\omega_2)$ bifurcation, 227
 - reversible $0(i\omega)$ bifurcation, 195
 - reversible 00^2 bifurcation, 193
 - reversible 0^20^2 bifurcation, 234
 - reversible 0^{2+} bifurcation, 160, 166
 - reversible $0^{2+}(i\omega)$ bifurcation, 199, 205
 - reversible 0^{2-} bifurcation, 173
 - reversible $0^{2-}(i\omega)$ bifurcation, 210
 - reversible 0^{3+} bifurcation, 181, 310
 - reversible 0^{3-} bifurcation, 192
 - reversible 0^{4+} bifurcation, 229
 - reversible systems, 118, 119
 - Takens–Bogdanov, 134
 - theorem, 94, 110, 117, 119, 144, 152, 153
 - time-periodic, 144, 147, 149
- ODE, 4
- operator
 - adjoint, 283
 - bounded, 279
 - closed, 280
 - compact, 282
 - Fredholm, 284
 - graph, 279
 - homological, 96
 - kernel, 30, 279
 - linear, 279
 - Nemitsky, 291
 - norm, 280
 - range, 30, 279
 - resolvent, 280
 - resolvent set, 30, 280
 - self-adjoint, 283
 - spectrum, 30
 - with compact resolvent, 282
- orbit
 - group, 28
 - heteroclinic, 175
 - homoclinic, 164
 - periodic, 15, 163, 164, 202
 - quasi-periodic, 202
- PDE, 22
- Poincaré map, 15
- problem
 - Bénard–Rayleigh, 249
 - Couette–Taylor, 244
- reduction function, 34
- resolvent
 - operator, 280
 - set, 30, 280
- resonant
 - strongly, 141, 228

- weakly, 141, 228
- SHE, 70, 223
- Sobolev space, 284
- solution, 31
 - antisymmetric, 80
 - equilibrium, 27, 28, 55, 158, 164
 - heteroclinic, 175
 - homoclinic, 164
 - periodic, 15, 127, 163, 164, 202
 - quasi-periodic, 202
 - roll, 75
 - symmetric, 80
- spatial dynamics, 261, 265, 272
- spectral
 - decomposition, 281
 - projection, 281
- spectrum, 30, 280
 - central, 32
 - extended, 281
 - stable, 32
 - unstable, 32
- symmetry, 254, 261
 - $O(2)$, 24, 28, 73–75, 129, 217, 245, 247, 252, 257
 - $SO(2)$, 22, 24, 127, 234, 245
 - continuous, 55, 273
 - equivariance, 53, 64, 117, 207, 268
 - mirror, 5
 - reflection, 5, 207, 245, 252
 - reversibility, 54, 55, 60, 90, 118, 157, 207, 263, 273, 276
- system
 - Brusselator, 18, 81
 - Burgers, 63
 - equivariant, 53, 117
 - Euler, 259
 - Navier–Stokes, 239
 - reaction-diffusion, 270
 - reduced, 34, 49
 - reversible, 54, 118
 - van der Pol, 120
- water-wave
 - gravity-capillary, 259
 - three-dimensional, 264
 - two-dimensional, 261
- wave
 - almost planar, 271
 - breather, 275
 - convection roll, 253, 256, 257
 - Couette flow, 244
 - exterior corner, 271, 274
 - front, 278
 - helicoidal, 248
 - hexagon, 257
 - hexagonal convection cell, 256
 - hole, 271, 275
 - in lattices, 275
 - interior corner, 271, 274
 - modulated, 274
 - patchwork quilt, 257
 - planar, 271
 - regular triangle, 257
 - ribbon, 248
 - roll, 257
 - rotating, 24, 129, 133
 - solitary, 263, 275, 278
 - spiral, 248
 - standing, 133, 248
 - step, 271, 274
 - Taylor vortex flow, 247
 - traveling, 258
 - twisted vortex, 248
 - water-, 259
 - wavy inflow boundary, 248
 - wavy outflow boundary, 248
 - wavy vortex, 248